



**CEP Discussion Paper No 1055**

**June 2011**

**Recursive Contracts**

**Albert Marcet and Ramon Marimon**

## **Abstract**

We obtain a recursive formulation for a general class of contracting problems involving incentive constraints. These constraints make the corresponding maximization (*sup*) problems non recursive. Our approach consists of studying a recursive Lagrangian. Under standard general conditions, there is a recursive *saddle-point (infsup)* functional equation (analogous to a Bellman equation) that characterizes the recursive solution to the planner's problem and *forward-looking* constraints. Our approach has been applied to a large class of dynamic contractual problems, such as contracts with limited enforcement, optimal policy design with implementability constraints, and dynamic political economy models.

Keywords: Transactional relationships, contracts and reputation, recursive formulation, participation constraint

JEL Classifications: D80, L14

This paper was produced as part of the Centre's Macro Programme. The Centre for Economic Performance is financed by the Economic and Social Research Council.

## **Acknowledgements**

This is a substantially revised version of previously circulated papers with the same title (e.g. Marcet and Marimon 1998 and 1999). We would like to thank Fernando Alvarez, Truman Bewley, Edward Green, Robert Lucas, Andreu Mas-Colell, Fabrizio Perri, Edward Prescott, Victor Rios-Rull, Thomas Sargent, Robert Townsend and Jan Werner for comments on earlier developments of this work, all the graduate students who have struggled through a theory in progress and, in particular, Matthias Mesner and Nicola Pavoni for pointing out a problem overlooked in previous versions. Support from MCyT-MEyC of Spain, CIRIT, Generalitat de Catalunya and the hospitality of the Federal Reserve Bank of Minneapolis is acknowledged.

Albert Marcet is an Associate of the Macro Programme at the Centre for Economic Performance and Lecturer in Economics, London School of Economics. Ramon Marimon is Director of the Max Weber Postdoctoral Programme and Professor of Economics at the European University Institute (EUI). He is also Chairman of the Barcelona Graduate School of Economics.

Published by  
Centre for Economic Performance  
London School of Economics and Political Science  
Houghton Street  
London WC2A 2AE

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior permission in writing of the publisher nor be issued to the public or circulated in any form other than that in which it is published.

Requests for permission to reproduce any article or part of the Working Paper should be sent to the editor at the above address.

© A. Marcet and R. Marimon, submitted 2011

## 1 Introduction

Recursive methods have become a basic tool for the study of dynamic economic models. For example, Stokey, et al. (1989) and Ljungqvist and Sargent (2004) describe a large number of macroeconomic models that can be analysed using recursive methods. A main advantage of this approach is that it characterizes optimal decisions – at any time  $t$  – as time-invariant functions of a small set of state variables. In engineering systems, knowledge of the available technology and of the current state is enough to decide the optimal control, since current returns and the feasible set depend only on past and current predetermined variables. In this case the value of future states is assessed by the value function

and, under standard dynamic programming assumptions, the Bellman equation is satisfied and a standard recursive formulation is obtained.

However, one key assumption to obtain the Bellman equation is that future choices do not constrain the set of today's feasible choices. Unfortunately, this assumption does not hold in many interesting economic problems. For example, in contracting problems where agents are subject to intertemporal participation, or other intertemporal incentive constraints, the future development of the contract determines the feasible action today. Similarly, in models of optimal policy design agents' reactions to government policies are taken as constraints and, therefore, future actions limit the set of current feasible actions available to the government. Many dynamic games – for example, dynamic political-economy models – share the same feature that an agent's current feasible actions depend on functions of future actions.

In general, in the presence of forward-looking constraints – as in rational expectations models where agents commit to contracts subject to incentive constraints (e.g. commitment may be limited) – optimal plans, or contracts, do not satisfy the Bellman equation and the solution is not recursive in the standard sense. In this paper we provide an integrated approach for a recursive formulation of a large class of dynamic models with forward-looking constraints by reformulating them as equivalent recursive saddle-point problems.

Our approach has a wide range of applications. In fact, it has already proved to be useful in the study of very many models<sup>1</sup>. Just to mention a few examples: growth and business cycles with possible default (Marcet and Marimon (1992), Kehoe and Perri (2002), Cooley, *et al.* (2004)); social insurance (Atanasio and Rios-Rull (2000)); optimal fiscal and monetary policy design with incomplete markets (Aiyagari, Marcet, Sargent and Seppälä (2002), Svensson and Williams (2008)), and political-economy models (Acemoglu, Golosov and Tsyvinskii (2011)). For brevity, however, we do not present further applications here and limit the presentation of the theory to the case of full information.

We build on traditional tools of economic analysis such as duality theory of optimization, fixed point theory, and dynamic programming. We proceed in three steps. We first study the planner's problem with incentive constraints (**PP**) as an infinite-dimensional maximization problem, and we embed this problem in a more general class of planner's problems (**PP** <sub>$\mu$</sub> ); these problems are parameterized by the weight ( $\mu$ ) of a (Benthamite) social welfare function, which accounts for the functions appearing in the constraints with future controls (*forward-looking constraints*). The objective function of **PP** <sub>$\mu$</sub>  is similar to Pareto-optimal problems where  $\mu$  is the vector of weights given to the different infinitely-lived agents in the economy.

Second, we consider the Lagrangean which incorporates the *forward-looking constraints* of the first period, which defines our starting *saddle-point planner's problem* (**SPP** <sub>$\mu$</sub> ) and we prove a duality result between this saddle-point problem and the planner's problem (**PP** <sub>$\mu$</sub> ). This construction helps to characterize

---

<sup>1</sup>As we write this version google scholar reports that the working paper has been cited 290 times, many of these citations are applications of the method.

the ‘non-recursivity problem’ and provides a key step towards its resolution.

As it is well known, the solution of dynamic models with forward-looking constraints is, in general, time-inconsistent, in the following sense: if at some period  $t > 0$  the agent solves  $\mathbf{PP}_\mu$  for the whole future path given the state variables found at  $t$ , the agent will not choose the path that he had chosen in period zero (unless, of course, the forward-looking constraints are not binding, up to period  $t$ ). This ‘non-recursivity problem’ is at the root of the difficulties in expressing the optimal solution with a time-invariant policy function.

A key insight of our approach is to show that there is a *modified problem*  $\mathbf{PP}_{\mu'}$  such that if the agent reoptimizes this problem at  $t = 1$  for a certain  $\mu'$ , the solution from period  $t = 1$  onwards is the same that had been prescribed by  $\mathbf{PP}_\mu$  from the standpoint of period zero. The key is to choose the weights  $\mu'$  appropriately. We show that the appropriate  $\mu'$  is given by the Lagrange multipliers of  $\mathbf{SPP}_\mu$  in period zero. This procedure of sequentially connecting *saddle-point problems* is well defined and it is recursive when solutions are unique. The problem  $\mathbf{PP}_{\mu'}$  can be thought of as the ‘continuation problem’ that needs to be solved each period in order to implement the constrained-efficient solution. It supports our claim that the recursive formulation is obtained by introducing the vector  $\mu$ , summarizing the evolution of the Lagrange multipliers, as co-state variable in a time-invariant policy function. As a result, with our method it is easy to guarantee existence of the solution to  $\mathbf{PP}_{\mu'}$  for any  $\mu' \geq 0$ , making the practical implementation of this method not more complicated than standard dynamic programming problems.

Third, we extend dynamic programming theory to show that the sequence of *modified saddle-point problems* ( $\mathbf{SPP}_{\mu_t}$ ) satisfies a *saddle-point functional equation* ( $\mathbf{SPFE}$ ; a *saddle-point Bellman equation*) and, conversely, that policies obtained from solving the *saddle-point functional equation* ( $\mathbf{SPFE}$ ) provide a solution to the original  $\mathbf{SPP}_\mu$  and, therefore, to the  $\mathbf{PP}_\mu$  problem. This latter sufficiency result is very general; in particular, it does not rely on convexity assumptions. This is important because incentive constraints do not have a convex structure in many applications. However, this result is limited in that we assume (local) uniqueness of solutions. We discuss the role this assumption plays and, in particular, we show how our approach, and results, do not depend on this assumption.

In addition, we also show how standard dynamic programming results, based on a *contraction mapping theorem*, generalize to our *saddle-point functional equation* ( $\mathbf{SPFE}$ ). An immediate consequence of these results is that one can use standard computational techniques that have been used to solve dynamic programming problems – such as solution of first-order-conditions for a given recursive structure of the policy function, or value function iteration – to solve dynamic saddle-point problems. Not only the computational techniques needed but also our assumptions are standard in dynamic economic models.

Our approach is related to other existing approaches that study dynamic models with expectations constraints, in particular to the pioneering works of Abreu, Pearce and Stacchetti (1990), Green (1987) and Thomas and Worrall (1988), and the applications that have followed. We briefly discuss how these,

and other, works relate to ours in Section 6, after presenting the main body of the theory in Sections 4 and 5. Section 2 provides a basic introduction to our approach and Section 3 a couple of canonical examples (most proofs are contained in the Appendix).

## 2 Formulating contracts as recursive saddle-point problems

In this section we give an outline of our approach, leaving the technical details and proofs for sections 4 and 5. Our interest is to solve problems that have the following representation:

$$\mathbf{PP} \quad \sup_{\{a_t, x_t\}} \mathbb{E}_0 \sum_{t=0}^{\infty} \beta^t r(x_t, a_t, s_t) \quad (1)$$

$$\text{s.t. } x_{t+1} = \ell(x_t, a_t, s_{t+1}), \quad p(x_t, a_t, s_t) \geq 0, \quad t \geq 0 \quad (2)$$

$$\mathbb{E}_t \sum_{n=1}^{N_j+1} \beta^n h_0^j(x_{t+n}, a_{t+n}, s_{t+n}) + h_1^j(x_t, a_t, s_t) \geq 0, \quad j = 1, \dots, l, \quad t \geq 0 \quad (3)$$

$$x_0 = x, \quad s_0 = s,$$

$$a_t \text{ is measurable with respect to } (\dots, s_{t-1}, s_t).$$

where  $r, \ell, p, h_0, h_1$  are known functions,  $\beta, x, s$  known constants,  $\{s_t\}_{t=0}^{\infty}$  an exogenous stochastic Markov process,  $N_j = \infty$  for  $j = 0, \dots, k$ , and  $N_j = 0$  for  $j = k + 1, \dots, l$ .

Standard dynamic programming methods only consider constraints of form (2) (see, for example, Stokey, *et al.* (1989) and Cooley, (1995)). Constraints of form (3) are not a special case of (2), since they involve expected values of future variables<sup>2</sup>. We know from Kydland and Prescott (1977) that, under these constraints, the usual Bellman equation is not satisfied, the solution is *not*, in general, of the form  $a_t = f(x_t, s_t)$  for all  $t$ , and the whole history of past shocks  $s_t$  can matter for today's optimal decision. By letting  $N_j = \infty$  **PP** covers a large class of problems where discounted present values enter the implementability constraint. For example, long term contracts with *intertemporal participation constraints* take this form.<sup>3</sup> Alternatively, by letting  $N_j = 0$  **PP** covers problems where intertemporal reactions of agents must be taken into account. For example, dynamic Ramsey problems, where the government

<sup>2</sup>One might think that expressing (3) in the form  $v(x_t, s_t) - \psi(x_t, s_t) \geq 0$ , where  $v$  is the discounted sum  $\mathbb{E}_t \sum_{n=0}^{\infty} \beta^n h_0(x_{t+n}, a_{t+n}, s_{t+n}), x_t, a_t, s_t$  and  $\psi = h_1 - h_0$  converts (3) into (2). But this does not solve the problem since  $v$  is not known a priori.

<sup>3</sup>Combining (2) and (3) accounts for a broad class of constraints. For example, a nonlinear participation constraint of the form  $g(\mathbb{E}_t \sum_{n=0}^{\infty} \beta^n h(x_{t+n}, a_{t+n}, s_{t+n}), x_t, a_t, s_t) \geq 0$  can easily be incorporated in our framework with one constraint of the form (2),  $g(w_t, x_t, a_t, s_t) \geq 0$  (with control variables  $(w_t, a_t)$ ), and one of the form (3),  $\mathbb{E}_t \sum_{n=0}^{\infty} \beta^n h(x_{t+n}, a_{t+n}, s_{t+n}) = w_t$ .

chooses policy variables subject to optimal dynamic behavior by the agents in the economy, have this form<sup>4</sup>. Even though we focus on the two canonical cases  $N_j = \infty$  and  $N_j = 0$ , intermediate cases can be easily incorporated. It is then without loss of generality that we let  $N_j = \infty$ , for  $j = 0, \dots, k$ , and  $N_j = 0$  for  $j = k + 1, \dots, l$ .

A first step of our approach is to consider a more general class of problems, parameterized by  $\mu$ :

$$\mathbf{PP}_\mu \quad \sup_{\{a_t, x_t\}} \mathbb{E}_0 \sum_{j=0}^l \sum_{t=0}^{N_j} \beta^t \mu^j h_0^j(x_t, a_t, s_t)$$

$$\text{s.t. } x_{t+1} = \ell(x_t, a_t, s_{t+1}), \quad p(x_t, a_t, s_t) \geq 0, \quad (4)$$

$$\mathbb{E}_t \sum_{n=1}^{N_j+1} \beta^n h_0^j(x_{t+n}, a_{t+n}, s_{t+n}) + h_1^j(x_t, a_t, s_t) \geq 0, \quad t \geq 0, \quad (5)$$

$$x_0 = x, \quad s_0 = s \quad (6)$$

and  $a_t$  is measurable with respect to  $(\dots, s_{t-1}, s_t)$ .

The main difference with  $\mathbf{PP}$  is that in  $\mathbf{PP}_\mu$  we have incorporated the  $h_0^j$  functions of the *forward-looking* constraints (3) into the objective function. Also, the superindex  $j$  now starts from  $j = 0$ , with  $h_0^0$ , to account for the reward function of the original problem. More precisely, if we let  $h_0^0 = r$ , we set  $\mu = (1, 0, \dots, 0)$  and we choose a very large  $h_1^0$  to guarantee that (5) is never binding for  $j = 0$ ,  $\mathbf{PP}_\mu$  is the original  $\mathbf{PP}$ . Furthermore, it should also be noticed that the value function of this problem, when well defined – say,  $V_\mu(x, s)$  – is homogeneous of degree one in  $\mu$ ; a property that our approach exploits (and the reason for collecting in the objective function the original return function  $r$  of  $\mathbf{PP}$ , together with the forward-looking elements of the constraints).

Notice that  $\mathbf{PP}_\mu$  is an infinite-dimensional maximization problem which, under relatively standard assumptions, is guaranteed to have a solution *for arbitrary*  $\mu \geq 0$ . The solution is a *plan*<sup>5</sup>  $\mathbf{a} \equiv \{a_t\}_{t=0}$ , where  $a_t(\dots, s_{t-1}, s_t)$  is a state-contingent action (Proposition 1).

An intermediate step in our approach is to transform program  $\mathbf{PP}_\mu$  into a saddle-point problem in the following way. Consider writing the Lagrangean for  $\mathbf{PP}_\mu$  when a lagrange multiplier  $\gamma \in R^{l+1}$  is attached to the *forward-looking* constraints *only* in period  $t = 0$  and the remaining constraints for  $t > 0$  are left

<sup>4</sup>See Section 6 for references to related work using constraints of the form  $N_j = \infty$  and  $N_j = 0$ .

<sup>5</sup>We use the bold notation to denote sequences of measurable functions.

as constraints. The Lagrangean then is

$$L(\mathbf{a}, \gamma; \mu) = \mathbb{E}_0 \sum_{j=0}^l \sum_{t=0}^{N_j} \beta^t \mu^j h_0^j(x_t, a_t, s_t) + \sum_{j=0}^l \gamma^j \left( \mathbb{E}_0 \sum_{t=1}^{N_j+1} \beta^t h_0^j(x_t, a_t, s_t) + h_1^j(x_0, a_0, s_0) \right). \quad (7)$$

$$\quad (8)$$

If we find a saddle-point of  $L$  subject to (4) for all  $t \geq 0$  and (5) for all  $t \geq 1$ , the usual equivalences between this saddle-point and the optimal allocation of  $\mathbf{PP}_\mu$  can be exploited. Using simple algebra it is easy to show that  $L$  can be rewritten as the objective function in the following saddle point problem<sup>6</sup>:

$$\mathbf{SPP}_\mu \quad \inf_{\gamma \in \mathbb{R}_+^{l+1}} \sup_{\{a_t, x_t\}} \mu h_0(x_0, a_0, s_0) + \gamma h_1(x_0, a_0, s_0) + \beta \mathbb{E}_0 \sum_{j=0}^l \varphi^j(\mu, \gamma) \sum_{t=0}^{N_j} \beta^t h_0^j(x_{t+1}, a_{t+1}, s_{t+1}) \quad (9)$$

$$\text{s.t. } x_{t+1} = \ell(x_t, a_t, s_{t+1}), \quad p(x_t, a_t, s_t) \geq 0, \quad t \geq 0$$

$$\mathbb{E}_t \sum_{n=1}^{N_j+1} \beta^n h_0^j(x_{t+n}, a_{t+n}, s_{t+n}) + h_1^j(x_t, a_t, s_t) \geq 0, \quad j = 0, \dots, l, \quad t \geq 1$$

for initial conditions  $x_0 = 0$  and  $s_0 = s$ , where  $\varphi$  is defined as

$$\begin{aligned} \varphi^j(\mu, \gamma) &\equiv \mu^j + \gamma^j && \text{if } N_j = \infty, \text{ i.e. } j = 0, \dots, k \\ &\equiv \gamma^j && \text{if } N_j = 0, \text{ i.e. } j = k + 1, \dots, l. \end{aligned}$$

The usefulness of  $\mathbf{SPP}_\mu$  comes from the fact that its objective function has a very special form: the term inside the expectation in (9) is precisely the objective function of  $\mathbf{PP}_{\varphi(\mu, \gamma^*)}$  given the states  $(x_1^*, s_1)$ . This will allow us to show that if  $(\{a_t^*\}_{t=0}^\infty, \gamma^*)$  solves  $\mathbf{SPP}_\mu$  for initial conditions  $(x, s)$  then  $\{a_{t+1}^*\}_{t=0}^\infty$  solves  $\mathbf{PP}_{\varphi(\mu, \gamma^*)}$  given initial conditions  $(x_1^*, s_1)$ , where  $x_1^* = \ell(x_0, a_0^*, s_1)$ . That is, the continuation problem that needs to be solved in the next period is precisely a planner problem where the weights have been shifted according to  $\varphi$ .

We then show that, under fairly general conditions, solutions to  $\mathbf{PP}_\mu$  are solutions to  $\mathbf{SPP}_\mu$  (Theorem 1), and viceversa (Theorem 2). Also, the usual slackness conditions will guarantee that if  $(\{a_t^*\}, \gamma^*)$  solves  $\mathbf{SPP}_\mu$

$$\mathbb{E}_0 \sum_{j=0}^l \gamma^{j*} \left[ \sum_{t=1}^{N_j+1} \beta^t h_0^j(x_t^*, a_t^*, s_t) + h_1^j(x, a_0^*, s) \right] = 0, \quad (10)$$

so that the values achieved by the objective functions of  $\mathbf{SPP}_\mu$  and  $\mathbf{PP}_\mu$  coincide.

<sup>6</sup>We use the notation  $\mu h_0(x, a, s) \equiv \sum_{j=0}^l \mu^j h_0^j(x, a, s)$ .



If  $\mathbf{PP}_\mu$  were a standard dynamic programming problem (i.e. without (5)), then the following Bellman equation would be satisfied:

$$\begin{aligned} V_\mu(x, s) &= \sup_a \{ \mu h_0(x, a, s) + \beta \mathbb{E} [V_\mu(\ell(x, a, s'), s') \mid s] \} \\ \text{s.t. } & p(x, a, s) \geq 0. \end{aligned} \quad (11)$$

The reason that the Bellman equation holds is that in standard dynamic programming if  $\mathbf{PP}_\mu$  is reoptimized at period  $t = 1$  given initial conditions  $(x_1^*, s_1)$ , the reoptimization simply confirms the choice that had been previously made for  $t \geq 1$ . However, with forward-looking constraints (5) this Bellman equation is *not* satisfied, the reason being that if the problem is reoptimized at  $t = 1$  the choice will violate the forward looking constraint of period  $t = 0$ , if (5) is binding at  $t = 0$ . A central element of our approach is that, as suggested by the objective function of  $\mathbf{SPP}_\mu$ , if the solution is reoptimized in period  $t = 1$  with the new weights  $\mu' = \varphi(\mu, \gamma^*)$  – that is, if in period one the reoptimization is for the problem  $\mathbf{PP}_{\mu'}$ , – the result confirms the solution of the original problem  $\mathbf{PP}_\mu$ . This allows the construction of a recursive formulation of our original  $\mathbf{PP}_\mu$  problem where the value function with modified weights is included in the right-hand side of a functional ‘Bellman-like’ equation to capture the terms in (9). More specifically, we show that under fairly general assumptions solutions to  $\mathbf{SPP}_\mu$  obey a saddle-point functional equation (**SPFE**). More specifically, we look for functions  $W$  that satisfy the following

$$\begin{aligned} \mathbf{SPFE} \quad W(x, \mu, s) &= \inf_{\gamma \geq 0} \sup_a \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta \mathbb{E} [W(x', \mu', s') \mid s] \} \\ \text{s.t. } & x' = \ell(x, a, s'), p(x, a, s) \geq 0 \\ & \text{and } \mu' = \varphi(\mu, \gamma), \end{aligned}$$

and we show that this holds for  $W(x, \mu, s) = V_\mu(x, s)$  (Theorem 3).

We only consider problems where the infsup problem has unique optimal choices.<sup>7</sup> In this case optimal allocations and multipliers are uniquely determined and there is a *policy function*  $\psi$ ; i.e.  $(a^*, \gamma^*) = \psi(x, \mu, s)$  associated with a value function  $W$  satisfying **SPFE**. Finally we show that the following recursive formulation

$$\begin{aligned} (a_t^*, \gamma_t^*) &= \psi(x_t^*, \mu_t^*, s_t) \\ \mu_{t+1}^* &= \varphi(\mu_t^*, \gamma_t^*) \end{aligned}$$

gives the optimal policy we are seeking. More precisely, we first show that, given  $\{a_t^*, \gamma_t^*\}$  generated by  $\psi$  for initial conditions  $(x, \mu, s)$ ,  $(\{a_t^*\}_{t=0}^\infty, \gamma_0^*)$  solves  $\mathbf{SPP}_\mu$  in state  $(x, s)$ , that  $(\{a_t^*\}_{t=1}^\infty, \gamma_1^*)$  solves  $\mathbf{SPP}_{\mu_1^*}$  in state  $(x_1^*, s_1)$ , etc. (Theorem 4). As a result, the path  $\{a_t^*\}$  for  $\mu_0 = (1, 0, \dots, 0)$  is a solution to **PP**.

In this sense, the modified problem  $\mathbf{PP}_\mu$  is the ‘correct continuation problem’ to the planners’ problem: if  $\mu$  is properly updated, the solution can be found each period by re-optimizing  $\mathbf{PP}_{\mu_t^*}$ .

<sup>7</sup>We discuss this issue in more detail in Sections 4 and 5.

## 2.1 The Principle of Optimality with *forward-looking* constraints

Our central result is not a simple restatement of the standard dynamic programming *principle of optimality* to our saddle-point formulation<sup>8</sup>. This principle says (when there are no *forward-looking* constraints)<sup>9</sup>: if  $V_\mu$  satisfies the Bellman equation (11), evaluated at  $(x, s)$ , it is the (*sup*) value of  $\mathbf{PP}_\mu$ , when the initial state is  $(x, s)$ , and a sequence  $\{a_t^*\}_{t=1}^\infty$  solves  $\mathbf{PP}_\mu$  if and only if it satisfies:

$$\begin{aligned} V_\mu(x_t^*, s_t) &= \mu h_0(x_t^*, a_t^*, s) + \beta \mathbb{E} [V_\mu(x_{t+1}^*, s_{t+1}) | s_t] \\ x_{t+1}^* &= \ell(x_t^*, a_t^*, s_{t+1}), \quad x_t^* = x. \end{aligned}$$

In our context, under standard assumptions it is true that if  $\{a_t^*\}_{t=1}^\infty$  solves  $\mathbf{PP}_\mu$  when the initial state is  $(x, s)$  and attains the value  $V_\mu(x, s)$ , then  $W(x, \mu, s) \equiv V_\mu(x, s)$  and

$$W(x_t^*, \mu_t^*, s_t) = \mu_t^* h_0(x_t^*, a_t^*, s_t) + \gamma_t^* h_1(x_t^*, a_t^*, s_t) + \beta \mathbb{E} [W(x_{t+1}^*, \mu_{t+1}^*, s_{t+1}) | s_t] \quad (12)$$

$$\begin{aligned} x_{t+1}^* &= \ell(x_t^*, a_t^*, s_{t+1}), \quad x_t^* = x \\ \mu_{t+1}^* &= \varphi(\mu_t^*, \gamma_t^*), \quad \mu_0^* = (1, 0, \dots, 0), \end{aligned}$$

where  $\{\gamma_t^*\}_{t=1}^\infty$  is the sequence of Lagrange multipliers associated with the sequence of  $\mathbf{SPP}_\mu$  (Theorems 1 and 3).

However, the converse, sufficiency, theorem that if  $W(x, \mu, s)$  satisfies  $\mathbf{SPFE}$  and  $(\{a_t^*\}_{t=1}^\infty, \{\gamma_t^*\}_{t=1}^\infty)$  satisfies (12) then  $V_\mu(x, s) \equiv W(x, \mu, s)$  is the value of  $\mathbf{PP}_\mu$  and  $\{a_t^*\}_{t=1}^\infty$  solves  $\mathbf{PP}_\mu$  is true if, *in addition*,  $W(x, \mu, s) = \mu\omega(x, \mu, s)$  and

$$\omega_j(x_t^*, \mu_t^*, s_t) = h_0^j(x_t^*, a_t^*, s_t) + \beta \mathbb{E} [\omega_j(x_{t+1}^*, \mu_{t+1}^*, s_{t+1}) | s_t], \quad (13)$$

if  $j = 0, \dots, k$ , and

$$\omega_j(x_t^*, \mu_t^*, s_t) = h_0^j(x_t^*, \mu_t^*, s_t) \quad \text{if } j = k+1, \dots, l. \quad (14)$$

These recursive equations for the *forward-looking* constraints are needed to guarantee that these constraints are also satisfied in the original  $\mathbf{PP}_\mu$ . Notice that if  $W(x, \mu, s)$  is differentiable in  $\mu$  then, by *Euler's Theorem*,  $W(x, \mu, s) = \mu\omega(x, \mu, s)$  (where  $\omega_j \equiv \partial_{\mu^j} W$ ) and equations (13) and (14) follow from the *Envelope Theorem*. We show, and use, the fact that if  $\{a_t^*\}_{t=1}^\infty$  is unique (at least locally unique) then  $W(x, \mu, s)$  is differentiable in  $\mu$  and, therefore, we recover the *Principle of Optimality* for our saddle-point formulation (Theo-

<sup>8</sup>This subsection clarifies our approach and what is new with respect of our previous work; it can be skipped by the reader only interested in how our approach works and on our main results.

<sup>9</sup>See, for example, Stokey *et al* (1989).

rems 4 and 5), without having to impose equations (13) and (14) as ‘promise-keeping’ constraints<sup>10</sup>.

Before we turn to these results in Sections 4 and 5, in the next Section we show how our approach is implemented in a couple of canonical examples.

### 3 Two Examples

In this Section we illustrate our approach with two examples. In the first, there are only *intertemporal participation constraints* so it is a case when  $N_j = \infty$  (i.e.  $k = l$ ); in the second, there are only *intertemporal one-period (Euler) constraints* hence it is a case with  $N_j = 0$  (i.e.  $k = 0$ ). The first is similar to the model studied in Marcet and Marimon (1992), Kocherlakota (1996), Kehoe and Perri (2002), among others, and it is canonical of models with intertemporal default constraints; the second is based on the model studied by Aiyagari et al. (2002) and it is a canonical model with Euler constraints, as in Ramsey equilibria of optimal fiscal and monetary policy.

#### 3.1 Intertemporal participation constraints.

We consider, as an example, a model of a partnership, where several agents can share their individual risks and jointly invest in a project which can not be undertaken by single (or subgroups of) agents. Formally, there is a single good and  $J$  infinitely-lived consumers. Preferences of agent  $j$  are represented by  $E_0 \sum_{t=0}^{\infty} \beta^t u(c_t^j)$ ;  $u$  is assumed to be bounded, strictly concave and monotone, with  $u(0) = 0$ ;  $c$  represents individual consumption. Agent  $j$  receives an endowment of consumption good  $y_t^j$  at time  $t$  and, given a realization of the vector  $y_t$  agent  $j$  has an outside option that delivers total utility  $v_j^a(y_t)$  if he leaves the contract in period  $t$ , where  $v_j^a$  is some known function. It is often assumed that the outside option is the autarkic solution:  $v_j^a(y_t) = E \left[ \sum_{n=0}^{\infty} \beta^n u(y_{t+n}^j) \mid y_t^j \right]$ , which implicitly assumes that if agent  $j$  defaults in period  $t$  he is permanently excluded from the partnership and he does not have any claims on the production and the capital of the partnership in, and after, period  $t$ .

Total production is given by  $F(k, \theta)$ , and it can be split into consumption  $c$  and investment  $i$ . The stock of capital  $k$  depreciates at the rate  $\delta$ . The joint process  $\{\theta_t, y_t\}_{t=0}^{\infty}$  is assumed to be Markovian and the initial conditions  $(k_0, \theta_0, y_0)$  are given. The planner looks for pareto optimal allocations that ensure that no agent ever leaves the contract. Letting  $Y_t = \sum_{j=1}^J y_t^j > 0$  the planner’s problem

---

<sup>10</sup>In our previous work (Marcet and Marimon (1998, 1999)) we assumed uniqueness of solutions and used the fact that the contraction mapping guarantees the uniqueness of the value function. Messner and Pavoni (2004) showed how the *principle of optimality* could fail in our context when solutions are not unique. The missing element being the recursivity of the forward-looking constraints. The above statement of the *principle of optimality* for problems with forward-looking constraints and, correspondingly, our sufficiency theorems address this issue, which we further discuss in Section 6.

takes the form:

$$\begin{aligned}
\mathbf{PP} \quad & \max_{\{c_t, i_t\}} \mathbb{E}_0 \sum_{t=0}^{\infty} \beta^t \sum_{j=1}^J \alpha^j u(c_t^j) \\
& \text{s.t. } k_{t+1} = (1 - \delta)k_t + i_t, \\
& F(k_t, \theta_t) + Y_t - \left( \sum_{j=1}^J c_t^j + i_t \right) \geq 0, \text{ and} \\
& \mathbb{E}_t \sum_{n=0}^{\infty} \beta^n u(c_{t+n}^j) \geq v_j^a(y_t) \quad \text{for all } j, t \geq 0.
\end{aligned}$$

It is easy to map this planner's problem into our **PP** formulation. Let  $s \equiv (\theta, y)$ ;  $x \equiv k$ ;  $a \equiv (i, c)$ ;  $\ell(x, a, s) \equiv (1 - \delta)k + i$ ;  $p(x, a, s) \equiv F(k, \theta) + \sum_{j \in J} y^j - \left( \sum_{j \in J} c^j + i \right)$ ;  $r(x, a, s) \equiv \sum_{j=1}^J \alpha^j u(c^j)$ ,  $h_0^j(x, a, s) \equiv u(c^j)$ ,  $h_1^j(x, a, s) \equiv u(c^j) - v_j^a(y_t)$ ,  $j = 1, \dots, J$ . Problems **PP** $_{\mu}$  and **SPP** $_{\mu}$  are obtained mechanically insuring that (5) is not binding for  $j = 0$ .

Finally we obtain the recursive formulation that we are seeking. **SPFE** takes the form<sup>11</sup>

$$\begin{aligned}
W(k, \mu, y, \theta) &= \inf_{\gamma \geq 0} \sup_{c, i} \left\{ \sum_{j=1}^J (\mu^0 \alpha^j + \mu^j) u(c^j) + \gamma^j (u(c^j) - v_j^a(y)) \right. \\
&\quad \left. + \beta \mathbb{E} [W(k', \mu', y', \theta') | y, \theta] \right\} \\
\text{s.t. } k' &= (1 - \delta)k + i, \quad F(k, \theta) + \sum_{j=1}^J y^j - \left( \sum_{j=1}^J c^j + i \right) \geq 0 \\
&\text{and } \mu' = \mu + \gamma.
\end{aligned}$$

We know that  $W(k, \mu, y, \theta) = V_{\mu}(k, y, \theta)$  solves this functional equation. Letting  $\psi$  be the policy function associated with this functional equation and  $W$ , solutions to **PP** satisfy

$$\begin{aligned}
(c_t^*, i_t^*, \gamma_t^*) &= \psi(k_t^*, \mu_t^*, \theta_t, y_t) \\
\mu_{t+1}^* &= \mu_t^* + \gamma_t^*
\end{aligned}$$

with initial conditions  $(k_0, \mu_0, \theta_0, y_0)$ , where  $\mu_0 = (1, 0, \dots, 0)$ .

The planner would obtain the full commitment solution (subject to intertemporal participation constraints) from period  $t$  onwards if in period  $t = 1$  he solved **PP** $_{\mu_1^*}$  given initial conditions  $(k_1^*, \theta_1, y_1)$ , provided that the weights  $\alpha$  of the agents were adjusted according to  $\mu_1^*$ . Co-state variables  $\mu_t^*$  become the additional weight that the planner should assign to each agent above the initial weight  $\alpha^j$  if the planner reoptimizes in period one. The variable  $\mu_t$  is all that needs to be remembered from the past.

<sup>11</sup>Here we incorporate the knowledge that  $\gamma^{0*} = 0$ .

This recursive formulation allows easy computation of solutions using either first-order-conditions or value function iteration. It also helps in characterizing the solution to the optimal problem: the weights  $\mu_t^*$  evolve according to whether or not their participation constraints are binding. Every time that the participation constraint for an agent is binding, his weight is increased permanently by the amount of the corresponding Lagrange multiplier. An agent is induced not to default by increasing his consumption permanently, not only in the period where he is tempted to default, but smoothly over time.

Due to these changing weights relative marginal utilities across agents are not constant when participation constraints are binding, since the first-order-conditions imply

$$\frac{u'(c_t^i)}{u'(c_t^j)} = \frac{\alpha^j + \mu_{t+1}^j}{\alpha^i + \mu_{t+1}^i} \quad , \text{ for all } i, j \text{ and } t.$$

It follows that individual paths of consumption depend on individual histories (in particular, on past ‘temptations to default’) not just on the initial wealth distribution and the aggregate consumption path, as in the Arrow-Debreu competitive allocations, but that this dependence on the past is completely summarized by  $\mu_t$  (and, by homogeneity, the weights  $\alpha^j + \mu_{t+1}^j$  can be normalized to add up to one). This also shows that if enforcement constraints are never binding (e.g., punishments are severe enough) then  $\mu_t = \mu_0$  and we recover the “constancy of the marginal utility of expenditure”, and the “constant proportionality between individual consumptions,” given by  $u'(c_t^i)/u'(c_t^j) = \alpha^j/\alpha^i$ . In other words, the evolution of the co-state variables can also be interpreted as the evolution of the distribution of wealth. If intertemporal participation constraints are binding infinitely often there may be a non-degenerate distribution of consumption in the long-run; in contrast with an economy where intertemporal participation constraints cease to be binding, as in an economy with full enforcement.<sup>12</sup>

The evolution of the weights  $\mu$  also helps to characterize the decision for capital: the intertemporal Euler equation of **SPP** $_\mu$  is given by:

$$\mu_{t+1}^i u'(c_t^i) = \beta \text{E}_t [\mu_{t+2}^i u'(c_{t+1}^i) (F_{k_{t+1}} + (1 - \delta))] ,$$

That is, the ‘stochastic discount factor’,  $\beta u'(c_{t+1}^i)/u'(c_t^i)$  is distorted by  $(1 + \gamma_{t+1}^i/\mu_{t+1}^i)$ , a distortion which does not vanish unless the non-negative process  $\{\gamma_t^i\}$  converges to zero.

### 3.2 Intertemporal one-period constraints, A ramsey problem

We present an abridged version of the optimal taxation problem studied by Aiyagari et al. (2002). A representative consumer solves

<sup>12</sup>See, for example, Broer (2009) for a characterization of the non-degenerate stationary distribution of consumption, in a similar model with finite number of types and a continuum of agents of each type.

$$\begin{aligned}
& \max_{\{c_t, e_t, b_{t+1}\}} E_0 \sum_{t=0}^{\infty} \delta^t [u(c_t) + v(e_t)] \\
\text{s.t.} \quad & c_t + b_{t+1} p_t^b = e_t(1 - \tau_t) + b_t \\
& b_{t+1} \geq \underline{B}, \quad b_0 \text{ given,}
\end{aligned}$$

where  $c$  is consumption and  $e$  is effort (e.g. hours worked). The government must finance exogenous random expenditures  $g$  issuing debt and collecting taxes. Feasible allocations satisfy  $c_t + g_t = e_t$ . The budget of the government mirrors the budget of the representative agent. For convenience we assume the government can not get too much in debt due to a constraint  $b_{t+1} \geq \underline{B}$ . In a competitive equilibrium, the following intertemporal and intratemporal equations must be satisfied (provided  $b_{t+1} > \underline{B}$ ):

$$\begin{aligned}
p_t^b u'(c_t) &= \beta E_t u'(c_{t+1}) \\
-\frac{v'(e_t)}{u'(c_t)} &= 1 - \tau_t.
\end{aligned}$$

In a Ramsey equilibrium the government chooses sequences of taxes and debt that maximize the utility of the consumer subject to allocations being a competitive equilibrium allocations. Substituting the above equilibrium equations into the budget constraint of the consumer the Ramsey equilibrium can be found by solving

$$\begin{aligned}
\mathbf{PP} \quad & \max_{\{c_t, b_{t+1}\}} E_0 \sum_{t=0}^{\infty} \beta^t [u(c_t) + v(e_t)] \\
\text{s.t.} \quad & E_t [\beta b_{t+1} u'(c_{t+1})] = u'(c_t)(b_t - c_t) - e_t v'(e_t) \quad (15) \\
& b_{t+1} \geq \underline{B}, \quad b_0 \text{ given,}
\end{aligned}$$

where  $e_t = c_t + g_t$  is left implicit. This problem can be represented as a special case of **PP** by letting:  $s \equiv g$ ;  $x \equiv b$ ;  $a \equiv (c, b')$ ;  $p(x, a, s) \equiv b' - \underline{B}$ ;  $\ell(x, a, s') \equiv b'$ ;  $r(x, a, s) \equiv u(c) + v(e)$ ;  $h_0^1(x, a, s) \equiv \beta b u'(c)$ ;  $h_1^1(x, a, s) \equiv u'(c)(b - c) - e v'(e)$ . Problems **PP** $_{\mu}$  and **SPP** $_{\mu}$  are then easily defined.

Finally we obtain the recursive formulation that we are seeking. In its original notation, **SPFE** takes the form<sup>13</sup>

$$\begin{aligned}
W(b, \mu, g) &= \inf_{\gamma^1 \geq 0} \sup_{c, i} \{ \mu^0 [u(c_t) + v(e_t)] + \mu^1 b u'(c) \\
&\quad + \gamma^1 [u'(c)(b - c) - e v'(e)] \quad + \beta E [W(b', \mu', g') | g] \} \\
\text{s.t.} \quad & b' \geq \underline{B}, \mu^{0'} = \mu^0, \mu^{1'} = \gamma^1
\end{aligned}$$

<sup>13</sup>Here we incorporate the knowledge that  $\gamma^{0*} = 0$ .

Letting  $\psi$  be the policy function associated with this functional equation, efficient allocations satisfy

$$(c_t^*, b_{t+1}^*, \gamma_t^{1*}) = \psi(b_t^*, \mu_t^*, g_t)$$

for  $\mu_{t+1}^* = (1, \gamma_t^{1*})$  with initial conditions  $(b_0, \mu_0, g_0)$ , where  $\mu_0 = (1, 0)$ .

It is clear that in this case the only element of  $\mu$  that matters is the lagrange multiplier  $\gamma_{t-1}$ . The planner would obtain the full commitment solution from period  $t$  onwards if in period  $t = 1$  he solved  $\mathbf{PP}_{\mu_1^*}$  given initial conditions  $(b_1^*, g_1)$ ; that is, if the objective function would be modified to include the term  $\gamma_0^1 b_1 u'(c_1)$  in addition to the consumer's discounted utility from  $t = 1$  onwards. This term captures the commitment to enforcing the Euler equation (15) at  $t = 0$ .

This recursive formulation allows easy computation of solutions using either first-order-conditions or value function iteration. It also helps in characterizing the solution to the optimal problem. The first-order-conditions of the Ramsey problem imply that solutions satisfy

$$\mathbb{E}_t [(\gamma_t^1 - \gamma_{t+1}^1)u'(c_{t+1})] = 0. \quad (16)$$

As discussed in Aiyagari et al. (2002), with incomplete markets, this implies that  $\{\gamma_{1,t}^*\}$  is a non-negative submartingale. Lagrange multipliers modify the weight given to debt relative to the complete markets case. The optimal policy now can be understood as forcing the planner in each period to modify the deadweight loss of taxation with weight  $\gamma_{t-1}^*$ .

## 4 The relationship between $\mathbf{PP}_\mu$ , $\mathbf{SPP}_\mu$ , and $\mathbf{SPFE}$

This section proves the relationships between the initial maximization problem  $\mathbf{PP}_\mu$ , the saddle-point problem  $\mathbf{SPP}_\mu$  and the saddle-point functional equation  $\mathbf{SPFE}$  discussed in the previous Sections. We first describe the basic structure of the problems being considered.

### 4.1 Basic Structure

There exist an exogenous stochastic process  $\{s_t\}_{t=0}^\infty$ ,  $s_t \in S$ , defined on the probability space  $(S_\infty, \mathcal{S}, P)$ . As usual,  $s^t$  denotes a history  $(s_0, \dots, s_t) \in S_t$  and  $\mathcal{S}_t$  the  $\sigma$ -algebra of events of  $s^t$ ; while  $\{s_t\}_{t=0}^\infty \in S_\infty$ , with  $\mathcal{S}$  the corresponding  $\sigma$ -algebra. An action in period  $t$ , history  $s^t$ , is denoted by  $a_t(s^t)$ , where  $a_t(s^t) \in A \subset R^m$ ; when there is no confusion, it is simply denoted by  $a_t$ . Given  $s_t$  and the endogenous state  $x_t \in X \subset R^n$ , an action  $a_t$  is (technologically) feasible if  $p(x_t, a_t, s_t) \geq 0$ . If the latter feasibility condition is satisfied, the endogenous state evolves according to  $x_{t+1} = \ell(x_t, a_t, s_{t+1})$ . Plans,  $\mathbf{a} = \{a_t\}_{t=0}^\infty$ , are elements of  $\mathcal{A} = \{\mathbf{a} : \forall t \geq 0, a_t : S_t \rightarrow A \text{ and } a_t \in \mathcal{L}_\infty^m(S_t, \mathcal{S}_t, P), \}$ , where  $\mathcal{L}_\infty^m(S_t, \mathcal{S}_t, P)$  denotes the space of  $m$ -valued, essentially bounded,  $\mathcal{S}_t$ -measurable functions. The corresponding endogenous state variables are elements of  $\mathcal{X} = \{\mathbf{x} : \forall t \geq 0, x_t \in \mathcal{L}_\infty^n(S_t, \mathcal{S}_t, P)\}$ .

Given initial conditions  $(x, s)$ , a plan  $\mathbf{a} \in \mathcal{A}$  and the corresponding  $\mathbf{x} \in \mathcal{X}$ , the evaluation of such plan in  $\mathbf{PP}_\mu$  is given by

$$f_{(x,\mu,s)}(\mathbf{a}) = \mathbb{E}_0 \sum_{j=0}^k \sum_{t=0}^{N_j} \beta^t \mu^j h_0^j(x_t, a_t, s_t)$$

We can describe the forward-looking constraints by defining  $g : \mathcal{A} \rightarrow \mathcal{L}_\infty^{k+1}$  coordinatewise as

$$g(\mathbf{a})_t^j = \mathbb{E}_t \left[ \sum_{n=1}^{N_j+1} \beta^n h_0^j(x_{t+n}, a_{t+n}, s_{t+n}) \right] + h_1^j(x_t, a_t, s_t)$$

Given initial conditions  $(x, s)$ , the corresponding feasible set of plans is then

$$\begin{aligned} \mathcal{B}(x, s) = \{ \mathbf{a} \in \mathcal{A} : & p(x_t, a_t, s_t) \geq 0, g(\mathbf{a})_t \geq 0, \mathbf{x} \in \mathcal{X}, \\ & x_{t+1} = \ell(x_t, a_t, s_{t+1}) \text{ for all } t \geq 0, \text{ given } (x_0, s_0) = (x, s) \}. \end{aligned}$$

Then  $\mathbf{PP}_\mu$  can be written in compact form as

$$\mathbf{PP}_\mu \quad \sup_{\mathbf{a} \in \mathcal{B}(x,s)} f_{(x,\mu,s)}(\mathbf{a})$$

We denote solutions to this problem as  $\mathbf{a}^*$  and the corresponding sequence of state variables  $\mathbf{x}^*$ . When the solution exists we define the value function of  $\mathbf{PP}_\mu$  as

$$V_\mu(x, s) = f_{(x,\mu,s)}(\mathbf{a}^*) \quad (17)$$

Similarly, we can also write  $\mathbf{SPP}_\mu$  in a compact form, by defining

$$\begin{aligned} \mathcal{B}'(x, s) = \{ \mathbf{a} \in \mathcal{A} : & p(x_t, a_t, s_t) \geq 0, g(\mathbf{a})_{t+1} \geq 0; \mathbf{x} \in \mathcal{X} \\ & x_{t+1} = \ell(x_t, a_t, s_{t+1}) \text{ for all } t \geq 0, \text{ given } (x_0, s_0) = (x, s) \}. \end{aligned}$$

$$\mathbf{SPP}_\mu \quad \inf_{\gamma \in \mathbb{R}_+^k} \sup_{\mathbf{a} \in \mathcal{B}'(x,s)} \{ f_{(x,\mu,s)}(\mathbf{a}) + \gamma g(\mathbf{a})_0 \}$$

Note that  $\mathcal{B}'$  only differs from  $\mathcal{B}$  in that the forward-looking constraints in period zero  $g(\mathbf{a})_0 \geq 0$  are not included as a condition in the set  $\mathcal{B}'$ , instead these constraints form part of the objective function of  $\mathbf{SPP}_\mu$ .

## 4.2 Assumptions and existence of solutions to $\mathbf{PP}_\mu$

We consider the following set of assumptions:

- A1.**  $s_t$  takes values on a set  $S \subset \mathbb{R}^K$ .  $\{s_t\}_{t=0}^\infty$  is a Markovian stochastic process defined on the probability space  $(S_\infty, \mathcal{S}, P)$ .
- A2.** (a)  $X \subset \mathbb{R}^n$  and  $A$  is a closed subset of  $\mathbb{R}^m$ . (b) The functions  $p : X \times A \times S \rightarrow \mathbb{R}$  and  $\ell : X \times A \times S \rightarrow X$  are measurable and continuous.



- A3.** Given  $(x, s)$ , there exist constants  $B > 0$  and  $\varphi \in (0, \beta^{-1})$ , such that if  $p(x, a, s) \geq 0$  and  $x' = \ell(x, a, s')$ , then  $\|a\| \leq B \|x\|$  and  $\|x'\| \leq \varphi \|x\|$
- A4.** The functions  $h_i^j(\cdot, \cdot, s)$ ,  $i = 0, 1, j = 0, \dots, l$ , are continuous and uniformly bounded, and  $\beta \in (0, 1)$ .
- A5.** The function  $\ell(\cdot, \cdot, s)$  is linear and the function  $p(\cdot, \cdot, s)$  is concave.  $X$  and  $A$  are convex sets.
- A6.** The functions  $h_i^j(\cdot, \cdot, s)$ ,  $i = 0, 1, j = 0, \dots, l$ , are concave.
- A6s.** In addition to **A6**, the functions  $h_0^j(x, \cdot, s)$ ,  $j = 0, \dots, l$ , are strictly concave.
- A7.** For all  $(x, s)$ , there exists a program  $\{\tilde{a}_n\}_{n=0}^\infty$ , with initial conditions  $(x, s)$ , which satisfies the inequality constraints (4) and (5) with strict inequality.

Assumptions **A1** and **A2** are part of our basic structure, described in the previous sub-section. These assumptions, together with **A3-A4**, are standard and we treat them as our basic assumptions. Assumptions **A5-A7** are often made but they are not satisfied in some interesting models; however, these assumptions are only used in some of the results below. For example, the concavity assumptions **A5-A6** are not needed for many results, and assumption **A7** is a standard interiority assumption, only needed to guarantee the existence of Lagrange multipliers.

The following proposition gives sufficient conditions for a maximum to exist for any  $\mu$ . The aim is not to have the most general existence theorem<sup>14</sup>, but to stress that one can find fairly general conditions under which  $\mathbf{PP}_\mu$  has a solution for any  $\mu$ , which will be crucial in the discussion of how our approach compares with that of Abreu Pearce and Stachetti, since this ensures that the continuation problem (namely  $\mathbf{PP}_{\varphi(\mu, \gamma)}$ ) is well defined for any  $\gamma$ .

**Proposition 1.** Assume **A1-A6** and that the set of possible exogenous states  $S$  is countable. Fix  $(x, \mu, s) \in X \times R_+^{l+1} \times S$ . Assume there exists a feasible plan  $\tilde{\mathbf{a}} \in \mathcal{B}(x, s)$  such that  $f_{(\mu, x, s)}(\tilde{\mathbf{a}}) > -\infty$ . Then there exists a program  $\mathbf{a}^*$  which solves  $\mathbf{PP}_\mu$  with initial conditions  $x_0 = x, s_0 = s$ .

Furthermore, if **A6s** is also satisfied then the solution is (almost sure) unique.

**Proof:** See Appendix.

### 4.3 The relationship between $\mathbf{PP}_\mu$ and $\mathbf{SPP}_\mu$

The following result says that a solution to the maximum problem is also a solution to the saddle point problem. It follows from standard theory of constrained optimization in linear vector spaces (see, for example, Luenberger (1969, Section 8.3, Theorem 1 and Corollary 1)). As in the standard theory, convexity and

<sup>14</sup>For example, not requiring **A6** or the countability of  $S$ , which will require additional assumptions.

concavity assumptions (**A5 to A6**), as well as an interiority assumption (**A7**) are necessary in order to obtain the result.

**Theorem 1** ( $\mathbf{PP}_\mu \implies \mathbf{SPP}_\mu$ ). Assume **A1-A7** and fix  $\mu \in R_+^{l+1}$ . Let  $\mathbf{a}^*$  be a solution to  $\mathbf{PP}_\mu$  with initial conditions  $(x, s)$ . There exists a  $\gamma^* \in R_+^l$  such that  $(\mathbf{a}^*, \gamma^*)$  is a solution to  $\mathbf{SPP}_\mu$  with initial conditions  $(x, s)$ .

Furthermore, the value of  $\mathbf{SPP}_\mu$  is the same as the value of  $\mathbf{PP}_\mu$ , more precisely

$$V_\mu(x, s) = f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \quad (18)$$

**Proof:** It is an immediate application of Theorem 1 (8.3) in Luenberger (1969), p. 217.

The following is a theorem on the sufficiency of a saddle point for a maximum.

**Theorem 2** ( $\mathbf{SPP}_\mu \implies \mathbf{PP}_\mu$ ). Given any  $(x, \mu, s) \in X \times R_+^{l+1} \times S$ , let  $(\mathbf{a}^*, \gamma^*)$  be a solution to  $\mathbf{SPP}_\mu$  for initial conditions  $(x, s)$ . Then  $\mathbf{a}^*$  is a solution to  $\mathbf{PP}_\mu$  for initial conditions  $(x, s)$ .

Furthermore, the value of the two programs is the same and (18) holds.

Notice that Theorem 2 is a sufficiency theorem ‘almost free of assumptions.’ All that is needed is the basic structure of section 4.1 defining the corresponding infinite-dimensional optimization and saddle-point problems *together with* the assumption that a solution to  $\mathbf{SPP}_\mu$  exists. Once these conditions are satisfied assumptions **A2 to A7** are not needed.

**Proof:** The following proof is an adaptation, to  $\mathbf{SPP}_\mu$ , of a sufficiency theorem for Lagrangian saddle points (see, for example, Luenberger (1969), Theorem 8.4.2, p.221).

If  $(\mathbf{a}^*, \gamma^*)$  solves  $\mathbf{SPP}_\mu$ , minimality of  $\gamma^*$  implies that, for every  $\gamma \geq 0$ ,

$$(\gamma^* + \gamma) g(\mathbf{a}^*)_0 \geq \gamma^* g(\mathbf{a}^*)_0;$$

therefore,  $g(\mathbf{a}^*)_0 \geq 0$ , but since  $\mathbf{a}^* \in \mathcal{B}'(x, s)$ , it follows that  $\mathbf{a}^* \in \mathcal{B}(x, s)$ ; i.e.  $\mathbf{a}^*$  is a feasible program for  $\mathbf{PP}_\mu$ . Furthermore, the minimality of  $\gamma^*$  implies that

$$\gamma^* g(\mathbf{a}^*)_0 \leq 0 g(\mathbf{a}^*)_0 = 0$$

but since  $\gamma^* \geq 0$  and  $g(\mathbf{a}^*)_0 \geq 0$ , it follows that  $\gamma^* g(\mathbf{a}^*)_0 = 0$ . Now, suppose there exist  $\tilde{\mathbf{a}} \in \mathcal{B}(x, s)$  satisfying  $f_{(x, \mu, s)}(\tilde{\mathbf{a}}) > f_{(x, \mu, s)}(\mathbf{a}^*)$ , then, since  $\gamma^* g(\tilde{\mathbf{a}})_0 \geq 0$ , it must be that

$$f_{(x, \mu, s)}(\tilde{\mathbf{a}}) + \gamma^* g(\tilde{\mathbf{a}})_0 > f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0$$

which contradicts the maximality of  $\mathbf{a}^*$  for  $\mathbf{SPP}_\mu$ .

Finally, using  $\gamma^* g(\mathbf{a}^*)_0 = 0$ , we have  $f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 = V_\mu(x, s)$  ■

#### 4.4 The relationship between $\mathbf{SPP}_\mu$ , and $\mathbf{SPFE}$

Recall that a function  $W : X \times R_+^{l+1} \times S \rightarrow R$  satisfies  $\mathbf{SPFE}$  at  $(x, \mu, s)$  when

$$W(x, \mu, s) = \min_{\gamma \geq 0} \max_{a \in \mathcal{A}} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta \mathbb{E}[W(x', \mu', s') | s] \} \quad (19)$$

$$\text{s.t. } x' = \ell(x, a, s), \quad p(x, a, s) \geq 0 \quad (20)$$

$$\text{and } \mu' = \varphi(\mu, \gamma). \quad (21)$$

In (19) we have substituted inf sup by min max, implicitly assuming that a solution to the saddle point problem exists, in which case the value,  $W(x, \mu, s)$ , is uniquely determined<sup>15</sup>. In other words, the right side of  $\mathbf{SPFE}$  is well defined for all  $(x, \mu, s)$  and  $W$  for which a saddle point exists.

We say that  $W$  satisfies  $\mathbf{SPFE}$  if it satisfies  $\mathbf{SPFE}$  at *any possible state*  $(x, \mu, s) \in X \times R_+^{l+1} \times S$ . Given  $W$ , we define the *saddle-point policy correspondence* (*SP policy correspondence*)  $\Psi : X \times R_+^{l+1} \times S \rightarrow A \times R_+^{l+1}$  by

$$\begin{aligned} \Psi_W(x, \mu, s) = & \left\{ (a^*, \gamma^*) : a^* \in \arg \max_{a \in \mathcal{A}, x' \in X} \mu h_0(x, a, s) + \gamma^* h_1(x, a, s) + \beta \mathbb{E}[W(x', \mu^{*'}, s') | s] \right. \\ & \text{for } \mu^{*'} = \varphi(\mu, \gamma^*) \text{ and (20);} \\ & \gamma^* \in \arg \min_{\gamma \geq 0} \mu h_0(x, a^*, s) + \gamma h_1(x, a^*, s) + \beta \mathbb{E}[W(x^{*'}, \mu', s') | s], \\ & \left. \text{for } x^{*'} = \ell(x, a^*, s) \text{ and (21)} \right\} \end{aligned}$$

If  $\Psi_W$  is single valued, we denote it by  $\psi_W$ , and we call it a *saddle-point policy function* (*SP policy function*).

We define the function  $W^*(x, \mu, s) \equiv V_\mu(x, s)$ . The following theorem says that  $W^*$  satisfies  $\mathbf{SPFE}$ .

**Theorem 3 ( $\mathbf{SPP}_\mu \implies \mathbf{SPFE}$ ).** Assume that  $\mathbf{SPP}_\mu$  has a solution for any  $(x, \mu, s) \in X \times R_+^{l+1} \times S$ , then  $W^*$  satisfies  $\mathbf{SPFE}$ . Furthermore, letting  $(\mathbf{a}^*, \gamma^*)$  be a solution to  $\mathbf{SPP}_\mu$  at  $(x, s)$ , we have  $(a_0^*, \gamma^*) \in \Psi_{W^*}(x, \mu, s)$ .

As in Theorem 2, Theorem 3 is also a theorem ‘almost free of assumptions,’ once the underlying structure and the existence of a well defined solution to  $\mathbf{SPP}_\mu$  at all possible  $(x, \mu, s)$  is assumed.

**Proof:** By theorem 2, we have that whenever  $\mathbf{SPP}_\mu$  has a solution  $W^*$  is well defined. Then, we first prove that, for any given  $(x, \mu, s)$ , if  $(\mathbf{a}^*, \gamma^*)$  solves  $\mathbf{SPP}_\mu$  at  $(x, s)$  the following recursive equation is satisfied:

$$W^*(x, \mu, s) = \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbb{E}[W^*(x_1^*, \varphi(\mu, \gamma^*), s') | s] \quad (22)$$

<sup>15</sup>See Lemma 3A in Appendix B.

To prove  $\leq$  in (22) we write

$$\begin{aligned}
W^*(x, \mu, s) &= f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\
&= \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) \\
&\quad + \beta \mathbb{E} \left[ \sum_{j=0}^l \varphi^j(\mu, \gamma^*) \sum_{t=0}^{N_j} \beta^t h_0^j(x_{t+1}^*, a_{t+1}^*, s_{t+1}) \mid s \right] \\
&= \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbb{E} [f_{(x_1^*, \varphi(\mu, \gamma^*), s_1)}(\sigma \mathbf{a}^*) \mid s] \\
&\leq \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbb{E} [V_{\varphi(\mu, \gamma^*)}(x_1^*, s_1) \mid s] \\
&= \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbb{E} [W^*(x_1^*, \varphi(\mu, \gamma^*), s) \mid s]
\end{aligned}$$

where  $\sigma \mathbf{a}^*$  is the original optimal sequence shifted one period; formally, letting the shift operator  $\sigma : S^{t+1} \rightarrow S^t$  be given by  $\sigma(s^t) = (s_1, s_2, \dots, s_t)$ , we define the  $\mathcal{S}_{t+1}$ -measurable function  $\sigma a_t^*$  by  $\sigma a_t^*(s) \equiv a_{t+1}^*(s)$ . The first equality follows from the definition of  $W^*$  and because Theorem 2 guarantees (18), the second equality follows from the definition of  $f, g$  and simple algebra, the third equality follows from the definitions of  $f, \varphi$ , and  $\mathbf{a}^*$ . The weak inequality follows from the fact  $\mathbf{a}^*$  is a feasible solution for the problem  $\mathbf{PP}_{\varphi(\mu, \gamma^*)}$  with initial conditions  $(x_1^*, s_1)$  and that this program achieves  $V_{\varphi(\mu, \gamma^*)}(x_1^*, s_1)$  at its maximum, and the last equality follows from Theorem 2 and (18).

To show  $\geq$  (22) we construct a sequence  $\mathbf{a}^+$  that consists of the optimal choice for  $\mathbf{SPP}_\mu$  for initial conditions  $(x, s)$  in the initial period, but subsequently is followed by the optimal choices for  $\mathbf{PP}_{\varphi(\mu, \gamma^*(x, \mu, s))}$  for initial conditions  $(x_1^*, s_1)$ . To define  $\mathbf{a}^+$  formally we explicitly denote by  $(\mathbf{a}^*(x, \mu, s), \gamma^*(x, \mu, s))$  a solution to  $\mathbf{SPP}_\mu$  for given initial conditions  $(x, s)$  and we let

$$\begin{aligned}
a_0^+(x, \mu, s) &= a_0^*(x, \mu, s) \\
a_t^+(x, \mu, s) &= \sigma a_{t-1}^*(x_1^*, \varphi(\mu, \gamma^*(x, \mu, s), s), s_1)
\end{aligned}$$

for all  $(x, \mu, s)$  and  $t \geq 1$ . Also, we let  $\mathbf{x}^+$  be the corresponding sequence of state variables.

In what follows, we simplify again notation and we go back to denoting  $a_t^*(x, \mu, s)$  by  $a_t^*$ ,  $\gamma^*(x, \mu, s)$  by  $\gamma^*$ , and  $a_t^+(x, \mu, s)(s^t)$  by  $a_t^+$ ; then, we have:

$$\begin{aligned}
W^*(x, \mu, s) &= f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\
&\geq f_{(x, \mu, s)}(\mathbf{a}^+) + \gamma^* g(\mathbf{a}^+)_0 \\
&= \mu h_0(x, a_0^+, s) + \gamma^* h_1(x, a_0^+, s) \\
&\quad + \beta \mathbb{E} \left[ \sum_{j=0}^l \varphi(\mu, \gamma^*)^j \sum_{t=0}^{N_j} \beta^t h_0^j(x_{t+1}^+, a_{t+1}^+, s_{t+1}) \mid s \right] \\
&= \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) \\
&\quad + \beta \mathbb{E} [W^*(x_1^*, \varphi(\mu, \gamma^*), s_1) \mid s],
\end{aligned}$$

where the first equality has been argued before, the first inequality follows from the fact that  $\mathbf{a}^+(x, \mu, s)$  is a feasible allocation in  $\mathbf{SPP}_\mu$  for initial conditions  $(x, s)$  – but  $\mathbf{a}^*(x, \mu, s)$  is a solution of the max part to  $\mathbf{SPP}_\mu$  at  $(x, s)$ . The second equality just applies the definition of  $f$  and  $g$ , the last equality follows because  $\mathbf{a}^+$  is optimal for  $\mathbf{PP}_{\varphi(\mu, \gamma^*(x, \mu), s)}$  given initial conditions  $(x_1^*, s_1)$  from period 1 onwards and because Theorem 2 insures that (18) holds.

Notice that for this step of the proof it is crucial that we use  $\mathbf{SPP}_\mu$  in order to obtain a recursive formulation. The first inequality above only works because we are considering a saddle point problem. Indeed, the  $\mathbf{a}^+$  sequence (that reoptimizes in period  $t = 1$ ) is feasible for  $\mathbf{SPP}_\mu$  because this problem does not impose the forward looking constraints in  $t = 0$ . The sequence  $\mathbf{a}^+$  would not be feasible in the original problem  $\mathbf{PP}_\mu$ , because by reoptimizing at period  $t = 1$  the forward-looking constraints at  $t = 0$  would be typically violated.

This ends the proof of (22).

To show that  $W^*$  satisfies  $\mathbf{SPFE}$  we now prove that the right side of SPFE is well defined at  $W^*$  and that  $(a_0^*, \gamma^*)$  is a saddle point of the right side of (19) or, formally, that  $(a_0^*, \gamma^*) \in \Psi_{W^*}(x, \mu, s)$ .

We first prove that  $a_0^*$  solves the max part of the right side of SPFE. Given any  $\tilde{a} \in A, p(x, \tilde{a}, s) \geq 0$ , letting  $\tilde{a}_t^*(s^t) \equiv a_{t-1}^*(\ell(x, \tilde{a}, s'), \varphi(\mu, \gamma^*), s')(\sigma(s^t))$  for  $t \geq 1$ , the definition of  $\tilde{a}_t^*$ , and (18) give the first equality in

$$\begin{aligned}
& \mu h_0(x, \tilde{a}, s) + \gamma^* h_1(x, \tilde{a}, s) + \beta \mathbf{E} [W^*(\ell(x, \tilde{a}, s'), \varphi(\mu, \gamma^*), s') | s] \\
= & \mu h_0(x, \tilde{a}, s) + \gamma^* h_1(x, \tilde{a}, s) \\
& + \beta \mathbf{E} \left[ \sum_{j=0}^l \varphi(\mu, \gamma^*)^j \sum_{t=0}^{N_j} \beta^t h_0^j(\tilde{x}_{t+1}^*, \tilde{a}_{t+1}^*, s_{t+1}) | s \right] \\
= & f_{(x, \mu, s)}(\tilde{\mathbf{a}}^*) + \gamma^* g(\tilde{\mathbf{a}}^*)_0 \\
\leq & f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\
= & W^*(x, \mu, s)
\end{aligned}$$

the second equality follows by definition, and the inequality because  $(\mathbf{a}^*, \gamma^*)$  solves the max part of  $\mathbf{SPP}_\mu$  while the third equality follows from (18). Now we can combine this with (22) to obtain that for all feasible  $\tilde{a} \in A$

$$\begin{aligned}
& \mu h_0(x, \tilde{a}, s) + \gamma^* h_1(x, \tilde{a}, s) + \beta \mathbf{E} [W^*(x_1^*, \varphi(\mu, \gamma^*), s') | s] \\
\leq & \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbf{E} [W^*(x_1^*, \varphi(\mu, \gamma^*), s') | s]
\end{aligned}$$

implying that  $a_0^*$  solves the max part of the right side of the  $\mathbf{SPFE}$ .

A similar argument shows that  $\gamma^*$  solves the min part: for any  $\tilde{\gamma} \in R_+^{l+1}$  let now

$$\begin{aligned}
& \mu h_0(x, a_0^*, s) + \tilde{\gamma} h_1(x, a_0^*, s) + \beta \mathbb{E}[W^*(x_1^*, \varphi(\mu, \tilde{\gamma}), s') | s] \\
\geq & \mu h_0(x, a_0^*, s) + \tilde{\gamma} h_1(x, a_0^*, s) + \\
& + \beta \mathbb{E} \left[ \sum_{j=0}^l \varphi(\mu, \tilde{\gamma})^j \sum_{t=0}^{N_j} \beta^t h_0^j(x_{t+1}^*, a_{t+1}^*, s_{t+1}) | s \right] \\
= & f_{(x, \mu, s)}(\mathbf{a}^*) + \tilde{\gamma} g(\mathbf{a}^*)_0 \\
\geq & f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\
= & W^*(x, \mu, s) \\
= & \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbb{E}[W^*(\ell(x, a_0^*, s'), \varphi(\mu, \gamma^*), s') | s]
\end{aligned}$$

where the inequality follows from the fact that shifting the policies one period back of the plan  $\mathbf{a}^*$  is a feasible plan for the  $\mathbf{PP}_{\varphi(\mu, \tilde{\gamma})}$  problem with initial conditions  $(x_1^*, s')$  and that  $W^*(x_1^*, \varphi(\mu, \tilde{\gamma}), s')$  is the optimal value of  $\mathbf{PP}_{\varphi(\mu, \tilde{\gamma})}$ , the second inequality follows because  $(\mathbf{a}^*, \gamma^*)$  is a saddle point of  $\mathbf{SPP}_\mu$  and the equalities follow from definitions, Theorem 2 and (22).

Therefore  $\gamma^*$  solves the min part of the right side of **SPFE**.

Therefore  $(a_0^*, \gamma^*)$  is a saddle point of the right side of **SPFE**, this implies the first equality in

$$\begin{aligned}
& \min_{\gamma \geq 0} \max_{a \in \mathcal{A}} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta \mathbb{E}[W^*(x', \mu', s') | s] \} \\
= & \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) + \beta \mathbb{E}[W^*(x_1^*, \varphi(\mu, \gamma^*), s') | s] \\
= & W^*(x, \mu, s)
\end{aligned}$$

and the second equality comes, again, from (22). This proves that  $W^*$  satisfies **SPFE**. ■

The argument used in the proof of Theorem 3 can be iterated a finite number of times to show the underlying recursive structure of the  $\mathbf{PP}_\mu$  formulation. If  $\mathbf{PP}_\mu$  has a unique solution  $\{a_t^*\}_{t=0}^\infty$  at  $(x, s)$ , then by Theorem 1 there is a  $\mathbf{SPP}_\mu$  at  $(x, s)$  with solution  $(\{a_t^*\}_{t=0}^\infty, \gamma^*)$ , which in turn defines a  $\mathbf{PP}_{\varphi(\mu, \gamma^*)}$  problem. As it has been seen in the proof of Theorem 3,  $\{a_t^*\}_{t=1}^\infty$  solves  $\mathbf{PP}_{\varphi(\mu, \gamma^*)}$  at  $(\ell(x, a_0^*, s), s_1)$  and by Theorem 1 there is a  $\gamma_1^*$  such that  $(\{a_t^*\}_{t=1}^\infty, \gamma_1^*)$  solves  $\mathbf{SPP}_{\varphi(\mu, \gamma^*)}$  at  $(\ell(x, a_0^*, s), s_1)$ . In turn,  $\{a_t^*\}_{t=2}^\infty$  solves  $\mathbf{PP}_{\varphi^{(2)}(\mu, \gamma^*)}$  at  $(\ell^{(2)}(x, a_0^*, s), s_1)$  where  $\varphi^{(2)}(\mu, \gamma^*) \equiv \varphi(\varphi(\mu, \gamma^*), \gamma_1^*, s_1)$  and  $\ell^{(2)}(x, a_0^*, s) \equiv \ell(\ell(x, a_0^*, s), a_1^*, s_1)$ . Similarly, let  $\varphi^{(n+1)}(\mu, \gamma^*) \equiv \varphi(\varphi^{(n)}(\mu, \gamma^*), \gamma_n^*, s_n)$ , then by recursively applying the argument of the proof of Theorem 3 we obtain the following result.

**Corollary 3.1. (Recursivity of  $\mathbf{PP}_\mu$ ).** If  $\mathbf{PP}_\mu$  satisfies the assumptions of Theorem 1 and has a unique solution  $\{a_t^*\}_{t=0}^\infty$  at  $(x, s)$ , then, for any  $(t, x_t^*, s_t)$ ,  $\{a_{t+j}^*\}_{j=0}^\infty$  is the solution to  $\mathbf{PP}_{\varphi^{(t)}(\mu, \gamma^*)}$  at  $(x_t^*, s_t)$ , where  $\gamma^*$  is the minimizer of  $\mathbf{SPP}_\mu$  at  $(x, s)$ .

The value function has some interesting properties that we would like to emphasize. First, notice that

$$\begin{aligned} W^*(x, \mu, s) &= f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\ &= f_{(x, \mu, s)}(\mathbf{a}^*) \\ &= \mathbb{E}_0 \sum_{j=0}^l \sum_{t=0}^{N_j} \beta^t \mu^j h_0^j(x_t^*, a_t^*, s_t), \end{aligned}$$

therefore, if  $\{a_t^*\}_{t=0}^\infty$  at  $(x, s)$  is uniquely defined, then  $W^*$  has a unique representation

$$\begin{aligned} W^*(x, \mu, s) &= \sum_{j=0}^l \mu^j \omega_j^*(x, \mu, s) \\ &= \mu \omega^*(x, \mu, s) \end{aligned}$$

where, for  $j = 0, \dots, k$ ,  $\omega_j(x, \mu, s) \equiv \mathbb{E}_0 \sum_{t=0}^\infty \beta^t h_0^j(x_t^*, a_t^*, s_t)$ , and, for  $j = k + 1, \dots, l$ ,  $\omega_j(x, \mu, s) \equiv h_0^j(x_0^*, a_0^*, s_0)$ . Similarly, the value function of  $\mathbf{SPP}_{\varphi(\mu, \gamma^*)}$  at  $(x_1^*, s_1)$ ,  $x_1^* = \ell(x, a_0^*, s)$ , satisfies

$$W^*(x_1^*, \varphi(\mu, \gamma^*), s_1) \equiv \varphi(\mu, \gamma^*) \omega^*(x_1^*, \varphi(\mu, \gamma^*), s_1).$$

This representation not only has an interesting economic meaning – for example, as a ‘social welfare function,’ with varying weights, in problems with intertemporal participation constraints – but is also very convenient analytically. In particular, this representation shows<sup>16</sup> that  $W^*$  is *convex and homogenous of degree one* in  $\mu$ , with  $W^*(x, 0, s) = 0$ , for all  $(x, s)$ <sup>17</sup>. In addition, the following Corollary to Theorem 3 also shows that  $W^*$  satisfies what we call the *saddle-point inequality property SPI*. Lemmas 1 and 2 below show how these properties are extended to general  $W$  functions satisfying **SPFE**.

A function  $W(x, \mu, s) = \sum_{j=0}^l \mu^j \omega_j(x, \mu, s)$  satisfies the *saddle-point inequality property SPI* at  $(x, \mu, s)$  if and only if there exist  $(a^*, \gamma^*)$  satisfying

$$\begin{aligned} &\mu h_0(x, a^*, s) + \tilde{\gamma} h_1(x, a^*, s) + \beta \mathbb{E} [\varphi(\mu, \tilde{\gamma}) \omega(x', \varphi(\mu, \tilde{\gamma}), s') | s] \\ &\geq \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} [\varphi(\mu, \gamma^*) \omega(x', \varphi(\mu, \gamma^*), s') | s]. \end{aligned} \quad (23)$$

$$\geq \mu h_0(x, \tilde{a}, s) + \gamma^* h_1(x, \tilde{a}, s) + \beta \mathbb{E} [\varphi(\mu, \gamma^*) \omega(\tilde{x}', \varphi(\mu, \gamma^*), s') | s], \quad (24)$$

<sup>16</sup>See Lemma 2A in the Appendix B.

<sup>17</sup>A function which is convex, homogeneous of degree one and finite at 0, is also called a *sublinear function* (see Rockafellar, 1981, p.29).

for any  $\tilde{\gamma} \in R_+^{l+1}$  and  $(\tilde{a}, \tilde{x}')$  satisfying the technological constraints at  $(x, s)$ ; that is, in **SPI** the multiplier minimization is taken in relation to the optimal continuation values.

**Corollary 3.2. (SPP $_{\mu}$   $\implies$  SPI).** Let  $W^*(x, \mu, s) \equiv V_{\mu}(x, s)$  be the value of **SPP $_{\mu}$**  at  $(x, s)$ , for an arbitrary  $(x, \mu, s)$ , then  $W^*(x, \mu, s) = \sum_{j=0}^l \mu^j \omega_j^*(x, \mu, s)$  satisfies **SPI**.

**Proof:** We only need to show that (23) is satisfied, but this is immediate from the following identities

$$\begin{aligned} f_{(x, \mu, s)}(\mathbf{a}^*) &= \mu h_0(x, a_0^*, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j^*(x_1^*, \varphi(\mu, \gamma^*), s_1) \mid s \right] \\ \gamma g(\mathbf{a}^*)_0 &= \gamma [h_1(x, a_0^*, s) + \beta \mathbb{E} [\omega^*(x_1^*, \varphi(\mu, \gamma^*), s_1) \mid s]], \end{aligned}$$

and the definition of **SPP $_{\mu}$**  at  $(x, s)$ ; that is, for any  $\tilde{\gamma} \in R_+^{l+1}$ ,

$$\begin{aligned} &\mu h_0(x, a^*, s) + \tilde{\gamma} h_1(x, a^*, s) + \beta \mathbb{E} [\varphi(\mu, \tilde{\gamma}) \omega^*(x^{*'}, \varphi(\mu, \gamma^*), s') \mid s] \\ &= f_{(x, \mu, s)}(\mathbf{a}^*) + \tilde{\gamma} g(\mathbf{a}^*)_0 \\ &\geq f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\ &= \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} [\varphi(\mu, \gamma^*) \omega^*(x^{*'}, \varphi(\mu, \gamma^*), s') \mid s] \end{aligned}$$

■

We now show that, under fairly general conditions, programs satisfying **SPFE** are solutions to **SPP $_{\mu}$**  at  $(x, s)$ . More formally,

**Theorem 4 (SPFE  $\implies$  SPP $_{\mu}$ )** Assume  $W$ , satisfying **SPFE**, is continuous in  $(x, \mu)$  and convex and homogeneous of degree one in  $\mu$ . If the *SP policy correspondence*  $\Psi_W$ , associated with  $W$ , generates a solution  $(\mathbf{a}^*, \gamma^*)_{(x, \mu, s)}$ , where  $(\mathbf{a}^*)_{(x, \mu, s)}$  is uniquely determined, then  $(\mathbf{a}^*, \gamma^*)_{(x, \mu, s)}$  is also a solution to **SPP $_{\mu}$**  at  $(x, s)$ .

Notice that the assumptions on  $W$  are very general. In particular, if  $W(x, \mu, s)$  is the value function of **SPP $_{\mu}$**  at  $(x, s)$  (i.e.  $W(x, \mu, s) \equiv V_{\mu}(x, s)$ ) then (as Lemma 2A in Appendix B shows) it is convex and homogeneous of degree one in  $\mu$  and, if **A2** - **A5** are satisfied it is continuous and bounded in  $(x, \mu)$ . The only ‘stringent condition’ is that  $(\mathbf{a}^*)_{(x, \mu, s)}$  must be uniquely determined, which is the case when  $W$  is concave in  $x$  and **A6s** is satisfied ( see Corollary 4.1.).

Before proving these results, we show that, as we have seen for  $W^*$ , convex and homogeneous functions  $W$  satisfying **SPFE** have some interesting properties, which are used in the proof of Theorem 4. First, without loss of generality (see **F2** and **F3** in Appendix C), we can express the *recursive equation* (19) in the form

$$\begin{aligned} \mu \omega^d(x, \mu, s) &= \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) \\ &\quad + \beta \mathbb{E} [\varphi(\mu, \gamma) \omega^{d'}(x^{*'}, \varphi(\mu, \gamma^*), s') \mid s], \end{aligned} \quad (25)$$



where  $\mu\omega^d(x, \mu, s) = W(x, \mu, s)$ , and the vectors  $\omega^d$  and  $\omega^{d'}$  are (partial) directional derivatives, in  $\mu$ , of  $W(x, \mu, s)$  and  $W(x^*, \varphi(\mu, \gamma^*), s')$ , respectively. Therefore, the **SPFE saddle-point inequalities** take the form

$$\begin{aligned} & \mu h_0(x, a^*, s) + \tilde{\gamma} h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \tilde{\gamma}) \omega^{d'}(x^*, \varphi(\mu, \tilde{\gamma}), s') \mid s \right] \\ & \geq \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma^*) \omega^{d'}(x^*, \varphi(\mu, \gamma^*), s') \mid s \right] \end{aligned} \quad (26)$$

$$\geq \mu h_0(x, \tilde{a}, s) + \gamma^* h_1(x, \tilde{a}, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma^*) \omega^{d'}(\tilde{x}', \varphi(\mu, \gamma^*), s') \mid s \right], \quad (27)$$

for any  $\tilde{\gamma} \in R_+^{l+1}$  and  $(\tilde{a}, \tilde{x}')$  satisfying the technological constraints at  $(x, s)$ .

Second, as we show in Lemma 1, there is an equivalence between this **SPFE** property and the *saddle-point inequality property*, **SPI**, which substitutes (26) for

$$\begin{aligned} & \mu h_0(x, a^*, s) + \tilde{\gamma} h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \tilde{\gamma}) \omega^{d'}(x^*, \varphi(\mu, \gamma^*), s') \mid s \right] \\ & \geq \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma^*) \omega^{d'}(x^*, \varphi(\mu, \gamma^*), s') \mid s \right]. \end{aligned} \quad (28)$$

Third, as we show in Lemma 2, if in addition  $(\mathbf{a}^*, \boldsymbol{\gamma}^*)_{(x, \mu, s)}$  is uniquely determined, then  $W$  is differentiable in  $\mu$ . Alternatively, if  $W$  is not differentiable in  $\mu'$ , then different choices of  $\omega^{d'}$  can result in different solutions and the union of all these different solutions are the solutions to the saddle point problem, given by (28) and (27).

**Lemma 1 (SPI  $\iff$  SPFE).** If  $W(x, \cdot, s)$  is convex and homogeneous of degree one, then (26) is satisfied if and only if (28) is satisfied. Furthermore, the inequality (28) is satisfied if and only if the following conditions are satisfied, for  $j = 0, \dots, l$ ,

$$h_1^j(x, a_0^*, s) + \beta \mathbb{E} \left[ \omega_j^{d'}(x_1^*, \mu_1^*, s_1) \mid s \right] \geq 0 \quad (29)$$

$$\gamma^{*j} \left[ h_1^j(x, a_0^*, s) + \beta \mathbb{E} \left[ \omega_j^{d'}(x_1^*, \mu_1^*, s_1) \mid s \right] \right] = 0. \quad (30)$$

**Proof of Lemma 1:** That **SPI  $\implies$  SPFE** follows from **F4** (see Appendix C).

With respect to  $W(x^*, \varphi(\mu, \gamma), s')$ , **F4** takes the form:

$$\varphi(\mu, \tilde{\gamma}) \omega^{d'}(x^*, \varphi(\mu, \tilde{\gamma}), s') \geq \varphi(\mu, \tilde{\gamma}) \omega^{d'}(x^*, \varphi(\mu, \gamma^*), s');$$

therefore (28) together with this latter inequality results in the following inequalities, which show that (26) is satisfied whenever (28) is satisfied:

$$\begin{aligned} & \mu h_0(x, a^*, s) + \tilde{\gamma} h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \tilde{\gamma}) \omega^{d'}(x^*, \varphi(\mu, \tilde{\gamma}), s') \mid s \right] \\ & \geq \mu h_0(x, a^*, s) + \tilde{\gamma} h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \tilde{\gamma}) \omega^{d'}(x^*, \varphi(\mu, \gamma^*), s') \mid s \right] \\ & \geq \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma^*) \omega^{d'}(x^*, \varphi(\mu, \gamma^*), s') \mid s \right] \end{aligned}$$

To see that **SPFE**  $\implies$  **SPI**, let

$$G_{(x, a^*, s)}(\gamma, \mu) \equiv \mu h_0(x, a^*, s) + \gamma h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma) \omega^{d'}(x^{*'}, \varphi(\mu, \gamma), s') \mid s \right],$$

and

$$F_{(x, a^*, s)}(\gamma, \mu) \equiv \mu h_0(x, a^*, s) + \gamma h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma) \omega^{d'}(x^{*'}, \varphi(\mu, \gamma^*), s') \mid s \right],$$

Then (26) reduces to  $G_{(x, a^*, s)}(\gamma, \mu) \geq G_{(x, a^*, s)}(\gamma^*, \mu)$  and (28) to  $F_{(x, a^*, s)}(\gamma, \mu) \geq F_{(x, a^*, s)}(\gamma^*, \mu)$ . Since  $G_{(x, a^*, s)}(\gamma^*, \mu) = F_{(x, a^*, s)}(\gamma^*, \mu)$ , the above inequalities (??) show that, if  $f_{(x, a^*, s)}(\gamma^*, \mu) \in \partial_\gamma F_{(x, a^*, s)}(\gamma^*, \mu)$ , for all  $\gamma \geq 0$ , then

$$\begin{aligned} G_{(x, a^*, s)}(\gamma, \mu) - G_{(x, a^*, s)}(\gamma^*, \mu) &\geq F_{(x, a^*, s)}(\gamma, \mu) - F_{(x, a^*, s)}(\gamma^*, \mu) \\ &(\gamma - \gamma^*) f_{(x, a^*, s)}(\gamma^*, \mu); \end{aligned}$$

that is,  $f_{(x, a^*, s)}(\gamma^*, \mu) \in \partial_\gamma G_{(x, a^*, s)}(\gamma^*, \mu)$ .

Let  $g_{(x, a^*, s)}(\gamma^*, \mu)$  be an extreme point of  $\partial_\gamma G_{(x, a^*, s)}(\gamma^*, \mu)$ , since  $G_{(x, a^*, s)}(\gamma, \mu)$  is homogenous of degree one in  $\gamma$ , by **F2**  $\exists \gamma_k \longrightarrow \gamma^*$  with  $G$  differentiable at  $\gamma_k$  and  $\nabla G_{(x, a^*, s)}(y_k, \mu) \longrightarrow g_{(x, a^*, s)}(\gamma^*, \mu)$ . By homogeneity of degree zero of  $\omega^{d'}(x^{*'}, \mu', s')$  with respect to  $\mu'$

$$\nabla G_{(x, a^*, s)}(\gamma_k, \mu) = h_1(x, a^*(x, \mu, s), s) + \beta \mathbb{E} \left[ \omega^{d'}(x^{*'}(x, \mu, s), \varphi(\mu, \gamma_k), s') \mid s \right].$$

Given the differentiability of  $\nabla G_{(x, a^*, s)}(y_k, \mu)$  at  $\gamma_k$ , continuity<sup>18</sup> of  $\varphi$  and  $\omega^{d'}$ , implies that

$$g_{(x, a^*, s)}(\gamma^*, \mu) = h_1(x, a^*(x, \mu, s), s) + \beta \mathbb{E} \left[ \omega^{d'}(x^{*'}(x, \mu, s), \varphi(\mu, \gamma^*), s') \mid s \right],$$

and, therefore,  $g_{(x, a^*, s)}(\gamma^*, \mu) \in \partial_\gamma F_{(x, a^*, s)}(\gamma^*, \mu)$  – in fact, it is also an extreme point of  $\partial_\gamma F_{(x, a^*, s)}(\gamma^*, \mu)$ . This shows that  $\partial_\gamma F_{(x, a^*, s)}(\gamma^*, \mu) = \partial_\gamma G_{(x, a^*, s)}(\gamma^*, \mu)$ , which, in turn, implies the equivalence between (26) and (28).

Finally, the proof of the Kuhn-Tucker conditions is standard. First, necessity of (29) follows from the fact that  $\gamma^* \geq 0$  is finite, which will not be the case if, for some  $j = 0, \dots, l$ ,

$$h_1^j(x, a_0^*, s) + \beta \mathbb{E} \left[ \omega_j^{d'}(x_1^*, \mu_1^*, s_1) \mid s \right] < 0.$$

To see necessity of (30), let  $\gamma_{(i)}^{*j} = \gamma^{*j}$ , if  $j \neq i$ , and  $\gamma_{(i)}^{*i} = 0$ , then (28) results in:

$$\mu h_0(x, a^*, s) + \gamma_{(i)}^* h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma_{(i)}^*) \omega^{d'}(x^{*'}, \varphi(\mu, \gamma^*), s') \mid s \right]$$

<sup>18</sup>The continuity of  $\omega^{d'}$ , is given, for example, by Theorem 4F (& Corollary 4G) in Rockafellar (1981).

$$\geq \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} \left[ \varphi(\mu, \gamma^*) \omega^{d'}(x', \varphi(\mu, \gamma^*), s') \mid s \right],$$

which, together with (29), implies that

$$0 \geq \gamma^{*j} \left[ h_1^j(x, a_0^*, s) + \beta \mathbb{E} \left[ \omega_j^{d'}(x_1^*, \mu_1^*, s_1) \mid s \right] \right] \geq 0.$$

To see, that (29) and (30) imply (28), suppose they are satisfied and there exists a  $\tilde{\gamma} \geq 0$ , for which (28) it is not, then it must be that

$$\begin{aligned} & \tilde{\gamma} \left[ h_1(x, a_0^*, s) + \beta \mathbb{E} \left[ \omega^{d'}(x_1^*, \mu_1^*, s_1) \mid s \right] \right] \\ & < \gamma^* \left[ h_1(x, a_0^*, s) + \beta \mathbb{E} \left[ \omega^{d'}(x_1^*, \mu_1^*, s_1) \mid s \right] \right] = 0, \end{aligned}$$

which contradicts (29) ■

**Lemma 2.** If  $(\mathbf{a}^*, \gamma^*)_{(x, \mu, s)}$  is generated by  $\Psi_W(x, \mu, s)$  and  $(\mathbf{a}^*)_{(x, \mu, s)}$  is uniquely defined then  $W(x_t^*, \mu_t^*, s_t)$  is differentiable with respect to  $\mu_t^*$ ; for every  $(x_t^*, \mu_t^*, s_t)$ , with  $(x_t^*, \mu_t^*)$  realized by<sup>19</sup>  $(\mathbf{a}^*, \gamma^*)_{(x, \mu, s)}$ .

**Proof of Lemma 2:** By (30) the recursive equation (25) simplifies to

$$\mu \omega^d(x, \mu, s) = \mu h_0(x, a^*, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j^{d'}(x', \mu + \gamma^*), s' \mid s \right],$$

Assume, for the moment, that  $(\mathbf{a}^*, \gamma^*)_{(x, \mu, s)}$  is uniquely determined. By recursive iteration, it follows that

$$\begin{aligned} \mu \omega^d(x_0, \mu_0, s_0) &= \mu h_0(x_0, a_0^*, s_0) \\ &+ \beta \mathbb{E}_0 \left[ \sum_{j=0}^k \mu^j \left( h_0^j(x_1^*, a_1^*, s_1) + \beta \omega_j^{d''}(x_2^*, \mu + \gamma_0^* + \gamma_1^*), s_2 \right) \mid s_0 \right] \\ &= \mu h_0(x_0, a_0^*, s_0) + \beta \mathbb{E}_0 \left[ \sum_{j=0}^k \mu^j \sum_{t=1}^{\infty} \beta^t h_0^j(x_t^*, a_t^*, s_t) \mid s_0 \right]; \end{aligned}$$

therefore, uniqueness of  $(\mathbf{a}^*, \gamma^*)_{(x, \mu, s)}$  implies: *i*)  $\omega^d(x, \mu, s)$  is uniquely defined:  $\omega^d(x, \mu, s) = \omega(x, \mu, s) \equiv \nabla_{\mu} W(x, \mu, s)$ ; which, in turn, implies that  $W(x, \cdot, s)$  is differentiable, and *ii*)  $\omega_j(x, \mu, s) = \mathbb{E}_0 \sum_{t=0}^{\infty} \beta^t h_0^j(x_t^*, a^*(x_t^*, \mu_t^*, s_t), s_t)$ , for  $j = 0, \dots, k$  (with  $(x_0^*, \mu_0^*, s_0) \equiv (x, \mu, s)$ ,  $x_{t+1}^* = \ell(x_t^*, a^*(x_t^*, \mu_t^*, s_t), s_t)$ , and  $\mu_{t+1}^* = \mu^* + \gamma^*(x_t^*, \mu_t^*, s_t)$ ), and  $\omega_j(x, \mu, s) = h_0^j(x, a^*(x, \mu, s), s)$ , for  $j = k + 1 \dots l$

<sup>19</sup>That is,  $(x_0^*, \mu_0^*) \equiv (x, \mu)$ ,  $x_{t+1}^* = \ell(x_t^*, a_t^*, s_{t+1})$  and  $\mu_{t+1}^* = \varphi(\mu_t^*, \gamma_t^*)$ .

Given  $(\mathbf{a}^*)_{(x,\mu,s)}$ , suppose now  $(\mathbf{a}^*, \tilde{\gamma}^*)_{(x,\mu,s)}$  is also generated by  $\Psi_W(x, \mu, s)$ .

Both saddle-point paths must have the same value (see Lemma 3A in Appendix B); in particular, following the same recursive argument:

$$\begin{aligned} \mu\omega^d(x_0, \mu_0, s_0) &= \mu h_0(x_0, a_0^*, s_0) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j^{d'}(x^{*j}, \mu + \tilde{\gamma}^*, s') \mid s \right] \\ &= \mu h_0(x_0, a_0^*, s_0) + \beta \mathbb{E}_0 \left[ \sum_{j=0}^k \mu^j \sum_{t=1}^{\infty} \beta^t h_0^j(x_t^*, a_t^*, s_t) \mid s_0 \right], \end{aligned}$$

which proves the differentiability of  $W$  with respect to  $\mu$ , even when  $(\gamma^*)_{(x,\mu,s)}$  is not uniquely determined (i.e. there may be kinks in the Pareto frontier) ■

An immediate, and important, consequence of Lemma 2 is the following result:

**Corollary:** If  $(\mathbf{a}^*)_{(x,\mu,s)}$  is uniquely defined by  $\Psi_W(x, \mu, s)$ , from any initial condition  $(x, \mu, s)$ , then the following (recursive) equations are satisfied:

$$\omega_j(x, \mu, s) = h_0^j(x, a^*(x, \mu, s), s) + \beta \mathbb{E} [\omega_j(x^{*j}(x, \mu, s), \mu^{*j}(x, \mu, s), s') \mid s], \quad \text{if } j = 0, \dots, k, \quad \text{and} \quad (31)$$

$$\omega_j(x, \mu, s) = h_0^j(x, a^*(x, \mu, s), s) \quad \text{if } j = k+1, \dots, l. \quad (32)$$

Furthermore,  $(\mathbf{a}^*)_{(x,\mu,s)}$  is uniquely defined by  $\Psi_W(x, \mu, s)$  whenever  $W(\cdot, \mu, s)$  is concave and **A6s** is satisfied.

Notice that, in proving Lemma 2, uniqueness of the solution paths has implied uniqueness of the value function decomposition:  $W = \mu\omega$ . This unique decomposition has implied the recursive equations (31) and (31). Uniqueness of the value function decomposition is equivalent to the differentiability of the value function. In fact, once it has been established that the value function is differentiable, one can obtain equations (31) and (31) as a simple application of the *Envelope Theorem*. For example, equation (31) is just<sup>20</sup>:

$$\partial_j W(x, \mu, s) = h_0^j(x, a^*(x, \mu, s), s) + \beta \mathbb{E} [\partial_j W(x^{*j}(x, \mu, s), \mu^{*j}(x, \mu, s), s') \mid s].$$

We now turn to the proof of Theorem 4, where these recursive equations (31) and (31) play a key role.

**Proof (Theorem 4):** By Lemma 2, there is a unique representation  $W(x, \mu, s) = \mu\omega(x, \mu, s)$ . To see that solutions of **SPFE** satisfy the participation constraints of **SPP** $_{\mu}$ , we use the first-order-conditions (29) and (30), as well

---

<sup>20</sup>We use the standard notation  $\partial_j W(x, \mu, s) \equiv \frac{\partial W(x, \mu, s)}{\partial \mu_j}$ , and also  $\omega_j(x, \mu, s) \equiv \partial_j W(x, \mu, s)$

as the recursive equations of the *forward-looking* constraints (31) and (32) of the previous Corollary. As in the proof of Lemma 2, equation (31) can be iterated to obtain

$$\omega_j(x, \mu, s) = \mathbb{E}_0 \left[ \sum_{t=0}^{\infty} \beta^t h_0^j(x_t^*, a_t^*, s_t) | s \right], \text{ if } j = 0, \dots, k. \quad (33)$$

Following the same steps for any  $t > 0$  and state  $(x_t^*, \mu_t^*, s_t)$ , equation(32) and (33) together with the inequality (29) show that the intertemporal participation constraints in  $\mathbf{PP}_\mu$  – and therefore in  $\mathbf{SPP}_\mu$  – are satisfied; that is,

$$\mathbb{E}_t \sum_{n=1}^{N_j+1} \beta^n h_0^j(x_{t+n}^*, a_{t+n}^*, s_{t+n}) + h_1^j(x_t^*, a_t^*, s_t) \geq 0, ; t \geq 0, \quad j = 0, \dots, l \quad (34)$$

Now, to see that solutions of  $\mathbf{SPFE}$  are, in fact, solutions of  $\mathbf{SPP}_\mu$  we argue by contradiction. Suppose there exist a program  $\{\tilde{a}_t\}_{t=0}^{\infty}$ , and  $\{\tilde{x}_t\}_{t=0}^{\infty}$ ,  $\tilde{x}_0 = x, \tilde{x}_{t+1} = \ell(\tilde{x}_t, \tilde{a}_t, s_{t+1})$ , satisfying the constraints of  $\mathbf{SPP}_\mu$  with initial condition  $(x, s)$  and such that

$$\begin{aligned} & \mu h_0(x, \tilde{a}_0, s) + \gamma^* h_1(x, \tilde{a}_0, s) \\ & + \beta \mathbb{E} \left[ \sum_{j=0}^k (\mu^j + \gamma^{*j}) \sum_{n=1}^{\infty} \beta^n h_0^j(\tilde{x}_t, \tilde{a}_t, s_t) + \sum_{j=k+1}^l \gamma^{*j} h_0^j(\tilde{x}_1, \tilde{a}_1, s_1) | s \right] \\ & > \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) \\ & + \beta \mathbb{E} \left[ \sum_{j=0}^k (\mu^j + \gamma^{*j}) \sum_{n=1}^{\infty} \beta^n h_0^j(x_t^*, a_t^*, s_t) + \sum_{j=k+1}^l \gamma^{*j} h_0^j(x_1^*, a_1^*, s_1) | s \right]. \end{aligned} \quad (35)$$

The following string of equalities and inequalities, which we explain at the

end, contradict this inequality,

$$\begin{aligned} & \mu h_0(x, a_0^*, s) + \gamma_0^* h_1(x, a_0^*, s) \\ & + \beta \mathbb{E} \left[ \sum_{j=0}^k (\mu^j + \gamma^{*j}) \sum_{n=1}^{\infty} \beta^n h_0^j(x_t^*, a_t^*, s_t) + \sum_{j=k+1}^l \gamma^{*j} h_0^j(x_1^*, a_1^*, s_1) \mid s \right] \\ & = \mu h_0(x, a_0^*, s) + \gamma_0^* h_1(x, a_0^*, s) + \beta \mathbb{E} [\mu_1^* \omega(x_1^*, \mu_1^*, s_1) \mid s] \end{aligned} \quad (36)$$

$$\begin{aligned} & \geq \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) + \beta \mathbb{E} [\mu_1^* \omega(\tilde{x}_1, \mu_1^*, s_1) \mid s] \\ & = \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) \end{aligned} \quad (37)$$

$$\begin{aligned} & + \beta \mathbb{E} [\mu_1^* h_0(\tilde{x}_1, a^*(\tilde{x}_1, \mu_1^*, s_1), s_1) + \gamma^*(\tilde{x}_1, \mu_1^*, s_1) h_1(\tilde{x}_1, a^*(\tilde{x}_1, \mu_1^*, s_1), s_1)] \\ & \quad (38) \end{aligned}$$

$$\begin{aligned} & + \beta \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1) \omega(x^{*'}(\tilde{x}_1, \mu_1^*, s_1), \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1), s_2) \mid s] \\ & \geq \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) \\ & + \beta \mathbb{E} [\mu_1^* h_0(\tilde{x}_1, \tilde{a}_1, s_1) + \gamma^*(\tilde{x}_1, \mu_1^*, s_1) h_1(\tilde{x}_1, \tilde{a}_1, s_1)] \end{aligned} \quad (39)$$

$$\begin{aligned} & + \beta \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1) \omega(\tilde{x}_2, \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1), s_2) \mid s] \\ & \geq \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) \\ & + \beta \mathbb{E} [\mu_1^* h_0(\tilde{x}_1, \tilde{a}_1, s_1) + \gamma^*(\tilde{x}_1, \mu_1^*, s_1) h_1(\tilde{x}_1, \tilde{a}_1, s_1) + \beta \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1) \omega(\tilde{x}_2, \mu_1^*, s_2) \mid s] \\ & \quad (40) \end{aligned}$$

$$\begin{aligned} & \geq \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) \\ & + \beta \mathbb{E} [\mu_1^* [h_0(\tilde{x}_1, \tilde{a}_1, s_1) + \beta \omega(\tilde{x}_2, \mu_1^*, s_2)] \mid s] \end{aligned} \quad (41)$$

$$\begin{aligned} & = \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) + \beta \mathbb{E} [\mu_1^* h_0(\tilde{x}_1, \tilde{a}_1, s_1) \mid s] \\ & + \beta^2 \mathbb{E} [\mu_1^* h_0(\tilde{x}_2, a^*(\tilde{x}_2, \mu_1^*, s_2), s_2) + \gamma^*(\tilde{x}_2, \mu_1^*, s_2) h_1(\tilde{x}_2, a^*(\tilde{x}_2, \mu_1^*, s_2), s_2)] \\ & \quad (42) \end{aligned}$$

$$\begin{aligned} & + \beta \mu^{*'}(\tilde{x}_2, \mu_1^*, s_2) \omega(x^{*'}(\tilde{x}_2, \mu_1^*, s_2), \mu^{*'}(\tilde{x}_2, \mu_1^*, s_2), s_2) \mid s] \\ & \geq \mu h_0(x, \tilde{a}_0, s) + \gamma_0^* h_1(x, \tilde{a}_0, s) \\ & + \beta \mathbb{E} \left[ \sum_{j=0}^k (\mu^j + \gamma^{*j}) \sum_{t=1}^{\infty} \beta^t h_0^j(\tilde{x}_t, \tilde{a}_t, s_t) + \sum_{j=k+1}^l \gamma^{*j} h_0^j(\tilde{x}_1, \tilde{a}_1, s_1) \mid s \right] \end{aligned} \quad (43)$$

Notice that the first equality (36) is just uses the value function decomposition, the other two equalities (38) and (42) are simple expansions of the of the saddle-point value paths (i.e., of (25)) and in these expansions equations (31) and (31) play a key role. Inequalities (37) and (39) follow from the maximality property of **SPFE**. Inequalities (40) and (41) require explanation. Inequality (40) follows from one of the properties of convex and homogeneous of degree one functions (i.e. **F4**:  $\widehat{\mu} \omega(\widehat{\mu}) \geq \widehat{\mu} \omega(\mu)$ , see Appendix), given that (40) is simply  $\mu^{*'}(\tilde{x}_1, \mu_1^*, s_1) \omega(\tilde{x}_2, \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1)) \geq \mu^{*'}(\tilde{x}_1, \mu_1^*, s_1) \omega(\tilde{x}_2, \mu_1^*, s_2)$ . Inequality (41) follows from applying the slackness inequality (29), as well as equations (32) and (33) to the plan generated by **SPFE** in state  $(\tilde{x}_2, \mu_1^*, s_2)$  (i.e. to  $\{a_t^*(\tilde{x}_2, \mu_1^*, s_2)\}_{t=2}^{\infty}$ ); these inequalities are needed to show that such plan satisfies the corresponding **SPP** constraints (34); that is,  $\left[ h_1^j(\tilde{x}_1, \tilde{a}_1, s_1) + \beta \omega_j(\tilde{x}_2, \mu_1^*, s_2) \right] \geq$

0,  $j = 0, \dots, l$ . Finally, since the equality (42) is simply the equality (38) after one iteration, repeated iterations result in the last inequality (43), which contradicts (35).

It only remains to show that the inf part of **SPP** is also satisfied. Reasoning again by contradiction, suppose there exist a  $\tilde{\gamma} \geq 0$  such that

$$\begin{aligned}
& \mu h_0(x, a_0^*, s) + \tilde{\gamma} h_1(x, a_0^*, s) \\
& + \beta \mathbb{E} \left[ \sum_{j=0}^k (\mu^j + \tilde{\gamma}^j) \sum_{n=1}^{\infty} \beta^n h_0^j(x_t^*, a_t^*, s_t) + \sum_{j=k+1}^l \tilde{\gamma}^j h_0^j(x_1^*, a_1^*, s_1) \mid s \right] \\
& < \mu h_0(x, a_0^*, s) + \gamma^* h_1(x, a_0^*, s) \\
& + \beta \mathbb{E} \left[ \sum_{j=0}^k (\mu^j + \gamma^{*j}) \sum_{n=1}^{\infty} \beta^n h_0^j(x_t^*, a_t^*, s_t) + \sum_{j=k+1}^l \gamma^{*j} h_0^j(x_1^*, a_1^*, s_1) \mid s \right].
\end{aligned} \tag{44}$$

Using the value function decomposition representation, this inequality can also be expressed as

$$\begin{aligned}
& \tilde{\gamma} [h_1(x, a^*(x, \mu, s), s) + \beta \mathbb{E} [\omega(x^{*'}(x, \mu, s), \mu^{*'}(x, \mu, s), s') \mid s]] \\
& < \gamma^*(x, \mu, s) [h_1(x, a^*(x, \mu, s), s) + \beta \mathbb{E} [\omega_j(x^{*'}(x, \mu, s), \mu^{*'}(x, \mu, s), s') \mid s]],
\end{aligned}$$

but the first-order-conditions (29) and (30) require that (28) is satisfied, i.e.

$$\begin{aligned}
& \tilde{\gamma} [h_1(x, a^*(x, \mu, s), s) + \beta \mathbb{E} [\omega(x^{*'}(x, \mu, s), \mu^{*'}(x, \mu, s), s') \mid s]] \\
& \geq \gamma^*(x, \mu, s) [h_1(x, a^*(x, \mu, s), s) + \beta \mathbb{E} [\omega_j(x^{*'}(x, \mu, s), \mu^{*'}(x, \mu, s), s') \mid s]] = 0
\end{aligned}$$

which contradicts (44) ■

Corollary to Lemma 2 implies the following Corollary to Theorem 4:

**Corollary 4.1.** Assume  $W$ , satisfying **SPFE**, is continuous in  $(x, \mu)$ , convex and homogeneous of degree one in  $\mu$ , concave in  $x$  and that **A6s** is satisfied. If  $(\mathbf{a}^*, \boldsymbol{\gamma}^*)_{(x, \mu, s)}$  is generated by  $\Psi_W(x, \mu, s)$  then  $(\mathbf{a}^*, \boldsymbol{\gamma}^*)_{(x, \mu, s)}$  is also a solution to **SPP** $_{\mu}$  at  $(x, s)$ .

## 5 DSPP and the contraction mapping

In this Section we show how our main results –Theorems 3 and 4– can also be obtained by applying the *Contraction Mapping Theorem* to the *Dynamic Saddle-Point Problem*, corresponding to **SPFE**. This Section provides more general sufficient conditions to obtain a solution to the original problem **PP** $_{\mu}$  starting from **SPFE**. While these conditions are satisfied whenever the conditions of Theorem 4 are satisfied, they help to better understand the passage **SPFE**→**PP** $_{\mu}$  and, in particular, they show how the standard method of value function iteration extends to our *saddle-point problems* and, therefore, that computing solutions to our original **PP** $_{\mu}$  does not require special computational

techniques. Furthermore, it also shows the interest of using the  $W = \mu\omega$  representation in computing recursive contracts (i.e. taking  $\omega$  as the starting vector valued function) and how, in contrast with the ‘promise keeping’ approach to solving contractual problems, ‘promised values’ are not part of the constraints, but an outcome of the recursive contract<sup>21</sup>.

We first define some spaces of “value” functions

$$\begin{aligned} \mathcal{M}_b = \{ & W : X \times \mathcal{R}_+^{l+1} \times S \rightarrow \mathcal{R} \\ & i) W(\cdot, \cdot, s) \text{ is continuous, and } W(\cdot, \mu, s) \text{ bounded, when } \|\mu\| \leq 1, \\ & ii) W(x, \cdot, s) \text{ is convex and homogeneous of degree one} \} \end{aligned}$$

and

$$\begin{aligned} \mathcal{M}_{bc} = \{ & W \in \mathcal{M}_b \text{ and} \\ & iii) W(\cdot, \mu, s) \text{ is concave} \}. \end{aligned}$$

$\mathcal{M}_b$  is a space of continuous, bounded functions (in  $x$ ), and convex and homogenous of degree one (in  $\mu$ )<sup>22</sup>, while  $\mathcal{M}_{bc}$  is the subspace of concave functions (in  $x$ ). Both spaces are normed vector spaces with the norm

$$\|W\| = \sup \{|W(x, \mu, s)| : \|\mu\| \leq 1, x \in X, s \in S\}.$$

We show in Appendix D (Lemma 6A) that they are complete metric spaces; therefore, suitable spaces for the *Contraction Mapping Theorem*.

Since, whenever  $W$  satisfies (ii) it can be represented as  $W(x, \mu, s) = \mu\omega(x, \cdot, s)$  (see Lemma 4A), it is convenient to define the corresponding spaces of functions:

$$\begin{aligned} M_b = \{ & \omega : X \times \mathcal{R}_+^{l+1} \times S \rightarrow \mathcal{R}^{l+1} \text{ s.t., for } j = 0, \dots, l, \\ & i) \omega_j(\cdot, \cdot, s) \text{ is continuous, and } \omega_j(\cdot, \mu, s) \text{ bounded, when } \|\mu\| \leq 1 \\ & ii) \omega_j(x, \cdot, s) \text{ is convex and homogeneous of degree zero} \} \end{aligned}$$

and

$$\begin{aligned} M_{bc} = \{ & \omega \in M_b \text{ s.t., for } j = 0, \dots, l, \\ & iii) \omega_j(\cdot, \mu, s) \text{ is concave} \}. \end{aligned}$$

Notice that  $\omega \in M$  uniquely defines a function  $W \in \mathcal{M}$ , given by  $W \equiv \mu\omega$ , but  $W \in \mathcal{M}$  does not uniquely define a  $\mathcal{R}^{l+1}$  valued function  $\omega \in M$ ; it does when, in addition,  $W$  is differentiable in  $\mu$  (see Appendix C)<sup>23</sup>.

As we have seen in Section 4<sup>24</sup>, when  $W^*(x, \mu, s) = V_\mu(x, s)$  is the value of **SPP** $_\mu$ , with initial conditions  $(x, s)$ , then  $W^*(x, \mu, s) = \sum_{j=0}^l \mu^j \omega_j^*(x, \mu, s)$

<sup>21</sup>We further discuss the ‘promise keeping’ approach in Section 6.

<sup>22</sup>Without loss of generality, we could also require that  $W(x, 0, s) < \infty$  and then replace (ii) by  $W(x, \cdot, s)$  is *sublinear* (see footnote 12).

<sup>23</sup> $M$  denotes either  $M_b$  or  $M_{bc}$ .

<sup>24</sup>See also Lemma 2A in Appendix B.



with  $W^* \in \mathcal{M}_b$ , whenever **A2** - **A4** are satisfied (and  $W^* \in \mathcal{M}_{bc}$  if in addition **A5** - **A6** are satisfied); furthermore,  $\omega^* \in M$  is unique whenever  $(\mathbf{a}^*)_{(x,\mu,s)}$  is uniquely defined.

Given a function  $\omega \in M$ , and an initial condition  $(x, \mu, s)$ , we can define the following *Dynamic Saddle Point Problem*:

**DSPP**

$$\begin{aligned} & \inf_{\gamma \geq 0} \sup_a \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta \mathbb{E} [\mu' \omega(x', \mu', s') | s] \} \\ \text{s.t. } & x' = \ell(x, a, s), \quad p(x, a, s) \geq 0 \\ & \text{and } \mu' = \varphi(\mu, \gamma), \end{aligned}$$

To guarantee that this problem has well defined solutions we make an interiority assumption:

**A7b.** For any  $(x, s) \in X \times S$ , there exists an  $\tilde{a} \in A$ , satisfying  $p(x, a, s) > 0$ , such that, for any  $\mu' \in R_+^{l+1}$ ,  $\|\mu'\| < +\infty$ , and  $j = 0, \dots, l : h_1^j(x, \tilde{a}, s) + \beta \mathbb{E} [\omega^j(\ell(x, \tilde{a}, s'), \mu', s') | s] > 0$ .

Notice that **A7b** is satisfied, whenever **A7** is satisfied and  $\mu' \omega(\ell(x, \tilde{a}, s'), \mu', s')$  is the value function of **SPP** $_{(\ell(x, \tilde{a}, s'), \mu', s')}$ . In general, **A7b** is not a restrictive assumption in the class of possible value functions if the original problem has interior solutions. Nevertheless, an assumption, such as **A7b** is needed when one takes **DSPP** $_{(x,\mu,s)}$  as the starting problem. This is a relatively standard min max problem, except for the dependency of  $\omega$  on  $\varphi(\mu, \gamma)$ . The following proposition shows that it has a solution. Obviously, solutions to **DSPP** $_{(x,\mu,s)}$  satisfy **SPFE**. . An immediate consequence of **A7b**, is the following lemma:

**Lemma 3.** Assume **A4** and **A7b** and let  $\omega \in M_b$ . There exist a  $B > 0$ , such that if  $(a^*(x, \mu, s), \gamma^*(x, \mu, s))$  is a solution to **DSPP** at  $(x, \mu, s)$ , then  $\|\gamma^*(x, \mu, s)\| \leq B \|\mu\|$ .

**Proof:** Denote by  $(a^*, \gamma^*) \equiv (a^*(x, \mu, s), \gamma^*(x, \mu, s))$  the solution to **DSPP** at  $(x, \mu, s)$ , and let  $\tilde{a}$  be the interior solution of **A7b**, then

$$\begin{aligned} & \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta \mathbb{E} [\mu' \omega(\ell(x, a^*, s), \varphi(\mu, \gamma^*), s') | s] \\ &= \mu h_0(x, a^*, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j(\ell(x, a^*, s), \varphi(\mu, \gamma^*), s') | s \right] \\ &+ \gamma^* [h_1(x, a^*, s) + \beta \mathbb{E} [\omega(\ell(x, a^*, s), \varphi(\mu, \gamma^*), s') | s]] \\ &= \mu h_0(x, a^*, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j(\ell(x, a^*, s), \varphi(\mu, \gamma^*), s') | s \right] \\ &\geq \mu h_0(x, \tilde{a}, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j(\ell(x, \tilde{a}, s), \varphi(\mu, \gamma^*), s') | s \right] \\ &+ \gamma^* [h_1(x, \tilde{a}, s) + \beta \mathbb{E} [\omega(\ell(x, \tilde{a}, s), \varphi(\mu, \gamma^*), s') | s]] \end{aligned}$$

By assumption,

$$(\mu/\|\mu\|)h_0(x, a^*, s) + \beta \mathbf{E} \left[ \sum_{j=0}^k (\mu^j/\|\mu\|) \omega_j(\ell(x, a^*, s), \varphi((\mu/\|\mu\|), (\gamma^*/\|\mu\|)), s') \mid s \right]$$

is uniformly bounded ( **A4** and  $\omega \in M_b$  imply that there is uniform bound for the max value), while if  $(\gamma^{*j}/\|\mu\|) > 0$  then

$$(\gamma^{*j}/\|\mu\|) \left[ h_1^j(x, \tilde{a}, s) + \beta \mathbf{E} [\omega_j(\ell(x, \tilde{a}, s), \varphi((\mu/\|\mu\|), (\gamma^*/\|\mu\|)), s') \mid s] \right] > 0;$$

therefore, there must be a  $B > 0$  such that  $\|\gamma^*\| \leq B \|\mu\|$  ■

**Proposition 2.** Let  $\omega \in M_{bc}$  and assume **A1-A6** and **A7b**. There exists  $(a^*, \gamma^*)$  that solves **DSPP** $_{(x,\mu,s)}$ . Furthermore if **A6s** is assumed, then  $a^*(x, \mu, s)$  is uniquely determined.

**Proof:** It is a relatively standard proof of existence of an equilibrium, based on a fixed point argument; see Appendix D.

The following Corollary to Theorem 3, is a simple restatement of the theorem in terms of the *Dynamic Saddle Point Problem*

**Corollary 3.3.** (**SPP** $_{\mu}(x, s) \implies$  **DSPP** $_{(x,\mu,s)}$ ). Assume that **SPP** $_{\mu}$  at  $(x, s)$  has a solution  $(a^*, \gamma^*)$  with value  $V_{\mu}(x, s) = \sum_{j=0}^l \mu^j \omega_j^*(x, \mu, s)$ , then  $(a_0^*, \gamma^*)$  solves **DSPP** $_{(x,\mu,s)}$ .

When **DSPP** $_{(x,\mu,s)}$  has a solution, it defines a **SPFE** operator  $T^* : \mathcal{M} \longrightarrow \mathcal{M}$  given by

$$(T^*W)(x, \mu, s) = \min_{\gamma \geq 0} \max_a \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta \mathbf{E} [W(x', \mu', s') \mid s] \}$$

s.t.  $x' = \ell(x, a, s)$ ,  $p(x, a, s) \geq 0$   
and  $\mu' = \varphi(\mu, \gamma)$ ,

When  $W \equiv \mu\omega$ , with  $\omega \in M$ , and **DSPP** $_{(x,\mu,s)}$  uniquely defines the values  $h_0^j(x, a^*(x, \mu, s), s)$ ,  $j = 0, \dots, l$ , then  $T^*$  defines a mapping,  $T : M \longrightarrow M$ , given by

$$(T\omega_j)(x, \mu, s) = h_0^j(x, a^*(x, \mu, s), s) + \beta \mathbf{E} [\omega_j(x^*(x, \mu, s), \mu^*(x, \mu, s), s') \mid s], \quad (45)$$

if  $j = 0, \dots, k$ , and

$$(T\omega_j)(x, \mu, s) = h_0^j(x, a^*(x, \mu, s), s), \text{ if } j = k+1, \dots, l. \quad (46)$$

Two remarks are in order. First, as already said, notice that  $(T\omega_j)$  correspond to the ‘promise keeping’ approach to solving contractual problems but in our

approach ( $T\omega_j$ ) is not a constraint: it is an outcome. Second, a fixed point of  $T^*$  does not imply a fixed point of  $T$  when ‘the planner’ is indifferent to  $T$  reallocations (e.g.  $\mu_i = \mu_j$ ,  $\mu_i(T\omega_i) + \mu_j(T\omega_j) = \text{constant}$ ) resulting in multiple (indeterminate) continuation values (for  $i$  and  $j$ )<sup>25</sup>

**Proposition 3.** Assume **DSPP** has a solution, for any  $\omega \in M$  and  $(x, \mu, s)$ , then  $T^* : \mathcal{M} \rightarrow \mathcal{M}$  is a well defined contraction mapping. Let  $W^* = T^*(W^*)$  and  $W^* = \mu\omega^*$ , if in addition the solutions  $a^*(x, \mu, s)$  to **DSPP** are unique, then  $\omega^* = T(\omega^*)$  is unique.

**Proof:** The first part follows from showing that Blackwell’s sufficiency conditions for a contraction are satisfied for  $T^*$  (see Lemmas 7A to 10A in Appendix D), the second part from the definition of  $T$ .

Our last Theorem, Theorem 5, wraps up our sufficiency results and is, in fact, a Corollary to Theorem 4. It shows how, starting from a *Dynamic Saddle-Point Problem* and a corresponding well defined *Contraction Mapping* resulting in a unique value function, one obtains the solution to our original problem **PP** $_{\mu}$ . The previous Propositions 2 and 3 provide conditions guaranteeing that the assumptions of Theorem 5, regarding  $T$ , are satisfied.

**Theorem 5** (**DSPP** $_{(x, \mu, s)} \implies \mathbf{SPP}_{\mu}(x, s)$ ). Assume  $T : M \rightarrow M$  has a unique fixed point  $\omega^*$ . Then the *value function*  $W^*(\mu, x, s) = \mu\omega^*(x, \mu, s)$  is the value of **SPP** $_{\mu}$  at  $(x, s)$  and the solutions of **DSPP** define a *saddle-point correspondence*  $\Psi$ , such that if  $(\mathbf{a}^*, \boldsymbol{\gamma}^*)$  is generated by  $\Psi$  from  $(x, \mu, s)$ , then  $(\mathbf{a}^*, \boldsymbol{\gamma}^*)$  solves **SPP** $_{\mu}$  at  $(x, s)$  and  $\mathbf{a}^*$  is the unique solution to **PP** $_{\mu}$  at  $(x, s)$ .

**Proof:** By assumption **SPFE** is satisfied. The proof of Theorem 4 is based on having a unique representation  $W^*(\mu, x, s) = \mu\omega^*(x, \mu, s)$  which, in that proof, is given by Lemma 2. In Theorem 5 such a unique representation is assumed, which implicitly also means to assume that the values  $h_0^j(x, a^*(x, \mu, s), s) j = 0, \dots, l$  are uniquely determined, which in fact is all what is needed in the proof of Theorem 4.

## 6 Related work

Precedents of our approach can be found in Epple, Hansen and Roberds (1985), Sargent (1987) and Levine and Currie (1987), who introduced lagrange multipliers as co-state variables in linear-quadratic Ramsey problems. Similarly, recent studies of optimal monetary policy in sticky price models have included lagrange multipliers as co-states. Often, the reason given for including these past multipliers as co-states is the observation that past multipliers appear in

<sup>25</sup>Messner and Pavoni’s ‘counterexample’ is one of these cases of ‘flats in the Pareto frontier’, when one only considers the  $T^*$  map and not the  $T$  map; see Marimon, Messner and Pavoni (2010).

the first-order-conditions of the Ramsey problem. Our work provides a formal proof that, with standard assumptions, co-state past multipliers deliver the optimal solution in a general framework, encompassing a larger class of models with *forward-looking* constraints.

The pioneer work of Abreu, *et al.* (1990) -APS, from now on,- characterizing sub-game perfect equilibria, shows that past histories can be summarized in terms of promised utilities. Earlier related work was by Green (1987) and Thomas and Worrall (1988). This approach has been widely used in macroeconomics<sup>26</sup>. Some applications are by Kocherlakota (1995) in a model with participation constraints, Cronshaw and Luenberger (1994) in a dynamic game. Also, as in the earlier work Kydland and Prescott (1980), Chang (1998) and Phelan and Stacchetti (1999) study Ramsey equilibria using promised *marginal* utility as a state variable, and they note the analogy of their approach with APS's.

Both APS and our approach have in common that, starting from non-recursive problems, allow to obtain optimal solutions (obviously, the same solutions) where *forward-looking* constraints have a recursive structure. In relatively simple problems (e.g., convex problems of full information and low dimensionality, in terms of state variables and number of forward-looking constraints) the two approaches can be seen as mirrors of each other. Nevertheless, there is a conceptual difference which sets these two approaches further apart as more complex problems are analyzed: our state (including the co-state  $\mu$ ) is predetermined, while promised-utilities – as co-state variables – are not; furthermore, in the APS approach, in taking future promised-utilities as choice variables, the recursive structure of *forward-looking* constraints must be taken as ‘promise keeping’ constraints, while in our approach we obtain this recursive structure as a result (see, Subsection 2.1)<sup>27</sup>.

As it is well known promised utilities in the APS approach have to be restricted to lie in a set where the continuation problem is well defined, otherwise algorithmic computations break down. The set of feasible promised utilities is not known beforehand, it can only be characterized numerically, often leading to very complicated calculations. Whenever there are several natural state variables the set of feasible promised utilities is a function of the natural state variables and the problem of finding the set of feasible utilities is daunting. Considerable progress has been made either by improving algorithms or by re-defining the problem at hand<sup>28</sup>, but the issue of constraining promised utilities (or marginal utilities in a Ramsey problem) is always present.

One key advantage of our approach is that the continuation problem is given by  $\mathbf{PP}_{\mu'}$ , and it is easy to find standard assumptions guaranteeing that this problem has a solution for any co-state  $\mu' \geq 0$ ; in fact, the set of feasible co-states is known beforehand, it is simply the positive orthant. The difficulties

<sup>26</sup>Ljungqvist and Sargent (2000) provides an excellent introduction, and reference, to most of this recent work.

<sup>27</sup>We only provide a summary discussion of the contrast between both approaches, see Marcet (2008) for more details.

<sup>28</sup>See, for example, Abraham and Pavoni (2005) or Judd, Yeltekin and Conklin (2003).

associated with computing a set of co-states for which the continuation problem is well defined are absent.

A second advantage lies in the dimensionality of the decision vector. In the APS approach the planner has to decide at  $t$  the utility promised at all possible states in  $t + 1$ . If the underlying exogenous state variable can take  $N_s$  possible values the planner has to decide on at least  $N_s$  controls at  $t$ . Most applications of APS constrain themselves to assuming that the exogenous variable is binomial (say, it can be "high" or "low"), but if exogenous variables can take many values a high-dimensional decision vector has to be solved for. Again, there are ways of dealing with this, but it is no doubt an added difficulty. By contrast, in our approach the dimensionality of the decision variable is independent of  $N_s$ .

An additional issue is that the initial conditions for the state variables in our approach are given from the outset, namely  $\mu^0 = 1$  and  $\mu^j = 0$  for  $j \geq 1$ , while in the promised utility approach the promised utility in the first period has to be solved for separately. It is well known that for this it is needed that the Pareto frontier is downward slopping, otherwise the computations can become very cumbersome.<sup>29</sup>

Finally, an interesting -but not exclusive- feature of our approach is that the evolution of  $\mu$  often helps to directly characterize the behavior of the model. For example, in models with participation constraints the  $\mu$ 's allow to interpret the behavior of the model as changing the Pareto weights sequentially depending on how binding the participation constraints become. In Ramsey type models the behavior of the  $\mu$ 's is associated with the commitment technology and the role that budget constraints play in the objective function of the planner. We have discussed these interpretations in Section 3. Also, our approach facilitates to identify cases where despite the presence of forward-looking constraints the co-state variables do not need to be introduced in the model.<sup>30</sup>

The APS approach has some strengths over our approach. For example, it allows for the characterization of all feasible paths (not only the constrained-efficient) and it naturally applies to models with private information or models with multiple solutions. However, these initial advantages are also being overcome. Sleet and Yeltekin (2010) and Mele (2010) have extended our approach to address moral hazard problems. In problems with multiple (locally unique) solutions, it is also possible to find other feasible paths using our approach by changing the objective function. However, as the example of Messner and Pavoni (2004) shows, there are problems where optimal paths are bound to have a continuum of solutions (i.e. when the constrained-efficient Pareto frontier has flats). We have maintained the assumption of (local) uniqueness in this paper; nevertheless, Marimon, Messner and Pavoni (2011) have recently

---

<sup>29</sup>Even with two agents and a downward slopping Pareto frontier, as in Kocherlakota (1996), one may be interested in finding the efficient allocation that *ex-ante* gives the same utility to both agents. While this is trivial with our approach (just give the same initial weights in the **PP** problem), becomes very tricky with the APS approach since the 'right promise' must be made to determine the initial conditions.

<sup>30</sup>This can be the case, for example, in Cooley, Quadrini and Marimon (2004) and Anagnostopoulos, Cárceles-Poveda and Marcet (2011).

shown that there is a natural extension of our approach to solve problems with non-uniqueness <sup>31</sup>.

Many applications of our approach can be found in the literature, although it is beyond the scope of this paper to discuss them in detail. This seems to be testimony of the convenience of using our approach, specially in the presence of intertemporal participation constraints with natural state variables such as capital (as in Subsection 3.1) or first-order Euler equation constraints with bonds as natural state (as in Subsection 3.2).

Perhaps most interesting is that the approach here can be used as an intermediate step in solving models that go beyond the pure formulation of **PP**. For example, a second generation of models considers *endogenous* participation constraints, as in the non-market exclusion models of Cooley et al. (2004), Marimon and Quadrini (2011), and Ferrero and Marcet (2005). In these models the functions  $h$  that appear in the incentive constraints are endogenous, they depend on the optimal or equilibrium solution, and the approach of this paper is often used as an intermediate step, defining the underlying contracts. This allows the study of problems where the outside option is determined in equilibrium as in models of debt renegotiation and long-term contracts. Furthermore, the work of Debortoli and Nunes (2010) extends our approach to study models of partial commitment and political economy, and the work of Marimon and Rendhal (2011) extend is to study models where agents can behave strategically with respect to their participation constraints, as in dynamic bargaining problems.

## 7 Concluding remarks

We have shown that a large class of problems with implementability constraints can be analysed using an equivalent recursive saddle-point problem. This saddle-point problem obeys a saddle point functional equation, which is a version of the Bellman equation. This approach works for a very large class of models with incentive constraints: intertemporal enforcement constraint, intertemporal Euler equations in optimal policy policy or regulation design, etc. This means that a unified framework can be provided to analyse all these models. The key feature of our approach is that instead of having to write optimal contracts as history-dependent contracts one can write them as a stationary function of the standard state variables together with additional co-state variables. These co-state variables are – recursively – obtained from the Lagrange multipliers associated with the intertemporal incentive constraints, starting from pre-specified initial conditions. This simple representation also provides economic insight into the analysis of various contractual problems; for example, with intertemporal

---

<sup>31</sup>Marimon, Messner and Pavoni (2011) show how the results presented here can be applied when the co-state is extended with the ‘last non-negative multiplier’. Cole and Kubler (2010) also provide a solution for the non-uniqueness case. Their approach involves a mix of our approach and of the APS approach; they provide a solution for the two-agent case with intertemporal participation constraints.

participation constraints it shows how the (Bethamite) social planner changes the weights assigned to different agents in order to keep them within the social contract; in Ramsey optimal problems it shows the cost of commitment to the benevolent government.

We have provided here the first complete account of the basic theory of *recursive contracts*. Nevertheless, we had already expounded most of the elements of the theory in our previous work (in particular, Marcet and Marimon (1988 & 1999)), which has allowed others to built on it. Many applications are already found in the literature, showing the convenience of our approach, specially when natural state variables are present. Useful extensions are already available encompassing a larger set of problems than the ones considered here.

## 8 Authors' Affiliations

*Albert Marcet, London School of Economics and CEP,  
E-mail: a.marcet@lse.ac.uk*

*and*

*Ramon Marimon, European University Institute and Universitat Pompeu  
Fabra - BarcelonaGSE,*

*Via delle Fontanelle 20, I-50014 San Domenico di Fiesole, Italy;*

*Telephone: +39-055-4685809, Fax: +39-055-4685894,*

*E-mail: ramon.marimon@eui.eu*



## APPENDIX

### APPENDIX A (PROOF OF PROPOSITION 1)

The proof of Proposition 1 relies on the following result:

**Lemma 1A.** Assume **A1-A6** and that  $S$  is countable, then

- i)*  $\mathcal{B}(x, s)$  is non-empty, convex, bounded and  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  closed; therefore it is  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  compact
- ii)* Given  $d \in R$ , the set  $\{\mathbf{a} \in \mathcal{A} : f_{(x, \mu, s)}(\mathbf{a}) \geq d\}$  is convex and  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  closed

The proof of Lemma 1A builds on three theorems. First, the *Urysohn metrization theorem* stating that regular topological spaces with a countable base are metrizable<sup>32</sup>. Second, the *Mackey-Arens theorem* stating that different topologies consistent with the same duality share the same closed convex sets; in our case, the duality is  $(\mathcal{L}_\infty, \mathcal{L}_1)$  and the weakest and the strongest topology consistent with such duality; namely, the *weak-star*,  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  and the Mackey  $\tau(\mathcal{L}_\infty, \mathcal{L}_1)$ . Third, the *Alaoglu theorem* stating that norm bounded  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  closed subsets of  $\mathcal{L}_\infty$  are  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  compact<sup>33</sup>.

**Proof:**

Assumptions **A2**, and **A4 - A6** imply that  $\mathcal{B}(x, s)$  is convex, and closed under pointwise convergence. Since, by assumption  $S$  is countable, *Urysohn metrization theorem* guarantees that  $\mathcal{B}(x, s)$  is, in fact,  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  closed. Assumptions **A3** and **A4** imply that  $\mathcal{B}(x, s)$  is bounded in the  $\|\cdot\|_\infty^\beta$  norm as needed for compactness, according to the *Alaoglu theorem*.

Assumptions **A5** and **A6** imply that  $\mathcal{B}(x, s)$  and the upper contour sets  $\{\mathbf{a} \in \mathcal{A} : f_\mu(\mathbf{a}) \geq d\}$ , are convex and, together with the previous assumptions (**A2**, and **A4**), Mackey closed and, therefore,  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  closed<sup>34</sup> ■

**Proof of Proposition 1:** As in Bewley (1991), the central element of the proof follows from the *Hausdorff maximal principle* and an application of the *finite intersection property*<sup>35</sup>. Let  $\mathcal{P}_d = \{\mathbf{a} \in \mathcal{B}(x, s) : f_{(x, \mu, s)}(\mathbf{a}) \geq d\}$ ,

---

<sup>32</sup>See Dunford and Schwartz (1957) p. 24. in our case the metric we use is given by

$$\rho_\infty^\beta(\mathbf{a}, \mathbf{b}) = \sum_{n=0}^{\infty} \beta^n \sup_{s^n \in S^n} |a_n(s) - b_n(s)|.$$

<sup>33</sup>See Schaefer (1966) p. 130 and p. 84, respectively.

<sup>34</sup>See Bewley (1972) for a proof of the Mackey continuity expected utility, without assuming  $S$  being countable.

<sup>35</sup>See, Kelley (1955) p. 33-34. for the Hausdorff principle and the Minimal principle, and p. 136 for the theorem stating that "a set is compact if and only if every family of closed sets which has a the finite intersection property has a non-void intersection."

then by Lemma 1A ,  $\mathcal{P}_d$  is  $\sigma(\mathcal{L}_\infty, \mathcal{L}_1)$  closed. By the interiority assumption of Proposition 1, for  $d$  low enough, it is non-empty. In fact, we can consider the family of sets  $\{\mathcal{P}_d : d \in D\}$  for which  $\mathcal{P}_d \neq \emptyset$ , where  $D \subset R$ . The sets  $\mathcal{P}_d$  are ordered by inclusion; in fact, if  $d' > d$  then  $\mathcal{P}_{d'} \subset \mathcal{P}_d$  and every finite collection of them has a non-empty intersection (i.e.,  $\{\mathcal{P}_d : d \in D\}$  satisfies *the finite intersection property*), but then by compactness of  $\mathcal{B}(x, s)$  any family of subsets of  $\{\mathcal{P}_d : d \in D\}$  -say,  $\{\mathcal{P}_d : d \in B \subset D\}$ - has a non-empty intersection and, by inclusion, there is  $\mathcal{P}_{\hat{d}} = \cap \{\mathcal{P}_d : d \in B \subset D\} \neq \emptyset$ . In particular, there is  $\mathcal{P}_{d^*} = \cap \{\mathcal{P}_d : d \in D\} \neq \emptyset$  which -as the *the minimal principle* states- is a minimal member of the family  $\{\mathcal{P}_d : d \in D\}$ . It follows that if  $\mathbf{a}^* \in \mathcal{P}_{d^*}$  then  $f_{(x, \mu, s)}(\mathbf{a}^*) \geq f_{(x, \mu, s)}(\mathbf{a})$  for any  $\mathbf{a} \in \mathcal{B}(x, s)$ . Furthermore, if strictly concavity is assumed then  $\mathcal{P}_{d^*}$  must be a singleton, otherwise convex combinations of elements of  $\mathcal{P}_{d^*}$  will form a proper closed subset of  $\mathcal{P}_{d^*}$  contradicting its minimality ■

#### APPENDIX B (SOME PROPERTIES OF $W^*$ )

**Lemma 2A.** Let  $W^*(x, \mu, s) \equiv V_\mu(x, s)$  be the value of  $\mathbf{SPP}_\mu$  at  $(x, s)$ , for an arbitrary  $(x, \mu, s)$ , then

- i*)  $W^*(x, \cdot, s)$  is convex and homogeneous of degree one;
- ii*) if **A2- A4** are satisfied  $W^*(\cdot, \mu, s)$  is continuous and uniformly bounded, and
- iii*) if **A5** and **A6** are satisfied  $W^*(\cdot, \mu, s)$  is concave.

**Proof:** *i*) follows from the fact that, for any  $\lambda > 0$ ,  $f_{(x, \lambda\mu, s)}(\mathbf{a}) = \lambda f_{(x, \mu, s)}(\mathbf{a})$ . To see this, let  $(\gamma^*, \mathbf{a}^*)$  satisfy **SFPE**, i.e.

$$\begin{aligned} & f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma g(\mathbf{a}^*)_0 \\ & \geq f_{(x, \mu, s)}(\mathbf{a}^*) + \gamma^* g(\mathbf{a}^*)_0 \\ & \geq f_{(x, \mu, s)}(\mathbf{a}) + \gamma^* g(\mathbf{a})_0, \end{aligned}$$

for any  $\gamma \in R_+^{l+1}$  and  $\mathbf{a} \in \mathcal{B}'(x, s)$ . Then  $(\lambda\gamma^*, \mathbf{a}^*)$  satisfies

$$\begin{aligned} & f_{(x, \lambda\mu, s)}(\mathbf{a}^*) + \gamma g(\mathbf{a}^*)_0 \\ & \geq f_{(x, \lambda\mu, s)}(\mathbf{a}^*) + \lambda\gamma^* g(\mathbf{a}^*)_0 \\ & \geq f_{(x, \lambda\mu, s)}(\mathbf{a}) + \lambda\gamma^* g(\mathbf{a})_0, \end{aligned}$$

for any  $\gamma \in R_+^{l+1}$  and  $\mathbf{a} \in \mathcal{B}'(x, s)$ . *ii*) and *iii*) are straightforward; in particular, *ii*) follows from applying the Theorem of the Maximum (Stokey, Lucas and Prescott, 1989, Theorem 3.6) *iii*) follows from the fact that the constraint sets are convex and the objective function concave. ■

**Lemma 3A:** If the inf sup problem, **SPFE** at  $(x, \mu, s)$ , has a solution then the value of this solution is unique.

**Proof:** It is a standard argument: consider two solutions to the right side of **SPFE** at  $(x, \mu, s)$ ,  $(\tilde{a}, \tilde{\gamma})$  and  $(\hat{a}, \hat{\gamma})$ , then repeated application of the saddle-point condition implies:

$$\begin{aligned}
& \mu h_0(x, \tilde{a}, s) + \tilde{\gamma} h_1(x, \tilde{a}, s) + \beta \mathbb{E} [W^*(\ell(x, \tilde{a}, s'), \varphi(\mu, \tilde{\gamma}), s') | s] \\
& \geq \mu h_0(x, \hat{a}, s) + \tilde{\gamma} h_1(x, \hat{a}, s) + \beta \mathbb{E} [W^*(\ell(x, \hat{a}, s'), \varphi(\mu, \tilde{\gamma}), s') | s] \\
& \geq \mu h_0(x, \hat{a}, s) + \hat{\gamma} h_1(x, \hat{a}, s) + \beta \mathbb{E} [W^*(\ell(x, \hat{a}, s'), \varphi(\mu, \hat{\gamma}), s') | s] \\
& \geq \mu h_0(x, \tilde{a}, s) + \hat{\gamma} h_1(x, \tilde{a}, s) + \beta \mathbb{E} [W^*(\ell(x, \tilde{a}, s'), \varphi(\mu, \hat{\gamma}), s') | s] \\
& \geq \mu h_0(x, \tilde{a}, s) + \tilde{\gamma} h_1(x, \tilde{a}, s) + \beta \mathbb{E} [W^*(\ell(x, \tilde{a}, s'), \varphi(\mu, \tilde{\gamma}), s') | s]
\end{aligned}$$

therefore the value of the objective at both  $(\tilde{a}, \tilde{\gamma})$  and  $(\hat{a}, \hat{\gamma})$  coincides ■

#### APPENDIX C (SOME PROPERTIES OF CONVEX HOMOGENEOUS FUNCTIONS)

To simplify the exposition of these properties let  $F : R_+^m \rightarrow R$  continuous, convex and homogeneous of degree one. The *subgradient set* of  $F$  at  $y$ , denoted  $\partial F(y)$ , is given by

$$\partial F(y) = \{z \in R^m \mid F(y') \geq F(y) + (y' - y)z \text{ for all } y' \in R_+^m\}.$$

The following **facts**, regarding  $F$ , are used in proving Lemmas 1 and 2:

- F1.** If  $F$  is convex, then it is differentiable at  $y$  if, and only if,  $\partial F(y)$  consists of a single vector; i.e.  $\partial F(y) = \{\nabla F(y)\}$ , where  $\nabla F(y)$  is called the *gradient* of  $F$  at  $y$ .
- F2.** If  $F$  is convex and finite in a neighborhood of  $y$ , then  $\partial F(y)$  is the convex hull of the compact set
$$\{z \in R^m \mid \exists y_k \rightarrow y \text{ with } F \text{ differentiable at } y_k \text{ and } \nabla F(y_k) \rightarrow z\}.$$

**F3. Lemma 4A (Euler's formula).** If  $F$  is convex and homogeneous of degree one and  $z \in \partial F(y)$  then  $F(y) = yz$ .

**F4. Lemma 5A.** If  $F$  is convex and homogeneous of degree one, for any pair  $(f, \hat{f})$ , if  $f^d(y) \in \partial F(y)$  and  $f^d(\hat{y}) \in \partial F(\hat{y})$ , then  $\hat{y}f^d(\hat{y}) \geq \hat{y}f^d(y)$ .

**F1** is a basic result on differentiability of convex functions (see, Rockafellar, 1981, Theorem 4F, or 1970, Theorem 25.1). **F2** is a very convenient characterization of the subgradient set of a convex function (see Rockafellar, 1981, Theorem 4D, or 1970, Theorem 25.6). We now provide a proof of the last two facts.

**Proof of Lemma 4A:** Let  $z \in \partial F(y)$ , then for any  $\lambda > 0$ ,  $F(\lambda y) - F(y) \geq (\lambda y - y)z$ , and, by homogeneity of degree one:  $(\lambda - 1)F(y) \geq (\lambda - 1)yz$ . If  $\lambda > 1$  this weak inequality results in:  $F(y) \geq yz$ ; while if  $\lambda \in (0, 1)$  in  $F(y) \leq yz$  ■<sup>36</sup>

<sup>36</sup>Notice that this does not imply that  $F$  is linear, which requires that  $F(-y) = -F(y)$ .

**Proof of Lemma 5A:** To see **F4** notice that if  $f^d(y) \in \partial F(y)$  and  $f^d(\hat{y}) \in \partial F(\hat{y})$ , by convexity, homogeneity of degree one, and Euler's formula:  $F(\hat{y}) = \hat{y}f^d(\hat{y})$  and

$$\begin{aligned} F(\hat{y}) &\geq F(y) + (\hat{y} - y)f^d(y) \\ &= yf^d(y) + \hat{y}f^d(y) - yf^d(y) = \hat{y}f^d(y). \end{aligned}$$

#### APPENDIX D (PROOF OF PROPOSITIONS 2 AND 3)

**Proof of Proposition 2:** Given the assumptions of Proposition 2, for any  $(x, \mu, s)$ , and  $\gamma \in \mathcal{R}_+^{l+1}$ , let

$$\begin{aligned} F_{(x,\mu,s)}(\gamma) &= \arg \sup_a \left\{ \mu h_0(x, a, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j(x', \mu', s') \mid s \right] \right\} \\ \text{s.t. } x' &= \ell(x, a, s), p(x, a, s) \geq 0 \\ \text{and } \mu' &= \varphi(\mu, \gamma), \\ \text{and } h_1^j(x, a, s) + \beta \mathbb{E} [\omega^j(x', \mu', s') \mid s] &\geq 0, j = 0, \dots, l \end{aligned} \quad (47)$$

Since this is a standard maximization problem of a continuous function on a compact set, there is a solution  $a^*(x, \mu, s; \gamma) \in F_{(x,\mu,s)}(\gamma)$ . Furthermore, given that the constraint set is convex and has a non-empty interior (by **A2** and **A7b**), there is an associated multiplier vector; let  $\gamma^{*j}(x, \mu, s; \gamma)$  be the multiplier corresponding to (47), for  $j$ . In particular, by Lemma 3,  $\gamma^*(x, \mu, s; \gamma) \in G_{(x,\mu,s)}(a^*)$ , where

$$\begin{aligned} G_{(x,\mu,s)}(a) &= \arg \inf_{\{\gamma \geq 0 : \|\gamma\| \leq B\|\mu\|\}} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta \mathbb{E} [\varphi(\mu, \gamma) \omega(x', \varphi(\mu, \gamma), s') \mid s] \} \\ \text{s.t. } x' &= \ell(x, a, s), p(x, a, s) \geq 0 \end{aligned}$$

The rest of the proof is a trivial application of the *Theorem of the Maximum* (e.g. Stokey et al. (1989), p. 62) and of *Kakutani's Fixed Point Theorem* (e.g. Mas-Colell et al. (1995), p.953). First notice that  $\mu h_0(x, a, s) + \beta \mathbb{E} \left[ \sum_{j=0}^k \mu^j \omega_j(x', \mu', s') \mid s \right]$  is continuous in  $a$  and, by **A2**, **A4** and the definition of  $M_{bc}$ ,  $G_{(x,\mu,s)}^*(\cdot)$  is also continuous. Second, let  $A(x, s) \equiv \{a \in A : p(x, a, s) \geq 0\}$ , by **A2** and **A3**  $A(x, s)$  is compact and by **A5** is convex, while  $\Psi(\mu) \equiv \{\gamma \geq 0 : \|\gamma\| \leq B\|\mu\|\}$  is trivially compact and convex. Therefore,  $F_{(x,\mu,s)} : \Psi(\mu) \rightarrow A(x, s)$  and  $G_{(x,\mu,s)}^* : A(x, s) \rightarrow \Psi(\mu)$  are upper-hemicontinuous, non-empty and convex-valued correspondences, jointly mapping a convex and compact set,  $\Psi \times A(x, s)$ , in itself; by *Kakutani's Fixed Point Theorem* there is a fixed point  $(a^*, \gamma^*)$  which is a solution to **DSPP** $_{(x,\mu,s)}$ . Furthermore,  $F_{(x,\mu,s)}(\cdot)$  is a continuous function, when **A6s** is assumed  $\blacksquare$

**Lemma 6A.**  $\mathcal{M}$  is a nonempty complete metric space.

**Proof:** That it is non-empty is trivial. Except for the homogeneity property, that every Cauchy sequence  $\{W^n\} \in \mathcal{M}_{bc}$  converges to  $W \in \mathcal{M}_{bc}$  satisfying *i*), *iii*), and the convexity property *ii*), follows from standard arguments (see, for example, Stokey, et al. (1989), Theorem 3.1 and Lemma 9.5). To see that the homogeneity property is also satisfied, for any  $(x, \mu, s)$  and  $\lambda > 0$ ,

$$\begin{aligned} & |W(x, \lambda\mu, s) - \lambda W(x, \mu, s)| \\ = & |W(x, \lambda\mu, s) - W^n(x, \lambda\mu, s) + \lambda W^n(x, \mu, s) - \lambda W(x, \mu, s)| \\ \leq & |W(x, \lambda\mu, s) - W^n(x, \lambda\mu, s)| + \lambda |W^n(x, \mu, s) - W(x, \mu, s)| \\ \rightarrow & 0 \end{aligned}$$

■

**Lemma 7A.** Assume **A2** - **A6** and **A7b**. The operator  $T^*$  maps  $\mathcal{M}_{bc}$  into itself.

**Proof:** First, notice that by Proposition 2, given  $W \in \mathcal{M}_{bc}$ ,  $T^*W$  is well defined. The correspondences  $\Gamma : X \rightarrow X$  and  $\Phi : \mathcal{R}_+^{l+1} \rightarrow \mathcal{R}_+^{l+1}$ , defined by  $\Gamma(x)_{(\mu,s)} \equiv \{x' \in X : x' = \ell(x, a, s), p(x, a, s) \geq 0, \text{ for some } a \in A\}$  and

$$\Phi(\mu)_{(x,s)} \equiv \{\mu' \in \mathcal{R}_+^{l+1} : \mu' = \varphi(\mu, \gamma), \text{ for some } \gamma \in \mathcal{R}_+^{l+1}\},$$

are continuous and compact-valued (by **A2**, **A3** and **A5**, and the definition of  $\varphi$ , respectively) and, as in the proof of Proposition 2, by continuity of the objective function, it follows that  $T^*W(\cdot, \cdot, s)$  is continuous, and given **A3** and **A4**, and the boundedness condition on  $W$ , it follows that  $T^*W$  also satisfies *i*). To see that the homogeneity properties are satisfied, let  $(a^*, \gamma^*)$  satisfy

$$(T^*W)(x, \mu, s) = \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta EW(x^*, \mu^*, s')$$

then, for any  $\lambda > 0$

$$\lambda(T^*W)(x, \mu, s) = \lambda[\mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta EW(x^*, \mu^*, s')]$$

Furthermore,

$$\begin{aligned} & \lambda\mu h_0(x, a^*, s) + \lambda\gamma^* h_1(x, a^*, s) + \beta EW(x^*, \lambda\mu^*, s') \\ = & \lambda \left[ \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta EW(x^*, \mu^*, s') \right] \end{aligned}$$

To see that **SPFE** is satisfied, let  $\gamma \geq 0$ ,  $\mu' = \varphi(\lambda\mu, \gamma)$  and  $a \in A(x, s)$ ,  $x' = \ell(x, a, s')$ , then

$$\begin{aligned} & \lambda\mu h_0(x, a^*, s) + \gamma h_1(x, a^*, s) + \beta EW(x^*, \mu', s') \\ = & \lambda \left[ \mu h_0(x, a^*, s) + \gamma \lambda^{-1} h_1(x, a^*, s) + \beta EW(x^*, \mu' \lambda^{-1}, s') \right] \\ \geq & \lambda \left[ \mu h_0(x, a^*, s) + \gamma^* h_1(x, a^*, s) + \beta EW(x^*, \mu^*, s') \right] \\ \geq & \lambda \left[ \mu h_0(x, a, s) + \gamma^* h_1(x, a, s) + \beta EW(x', \mu^*, s') \right] \end{aligned}$$

It follows that,

$$\begin{aligned}(T^*W)(x, \lambda\mu, s) &= \lambda\mu h_0(x, a^*, s) + \lambda\gamma^* h_1(x, a^*, s) + \beta EW(x^*, \lambda\mu^*, s') \\ &= \lambda(T^*W)(x, \mu, s)\end{aligned}$$

Finally, since  $W \in \mathcal{M}_{bc}$ , it is straightforward to show that  $TW$  is concave in  $x$  (by **A5** and **A6**), and convex in  $\mu$  ■

**Lemma 7A (monotonicity)** Let  $\widehat{W} \in \mathcal{M}$  and  $\widetilde{W} \in \mathcal{M}$  be such that  $\widehat{W} \leq \widetilde{W}$ , then  $(T^*\widehat{W}) \leq (T^*\widetilde{W})$ .

**Proof** Fix  $(\mu, x, s)$ , then for any  $\mu'$  satisfying  $\mu' = \varphi(\mu, \gamma) \geq 0$ , for a given  $\gamma \geq 0$ ,

$$\begin{aligned}& \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta E \widehat{W}(\ell(x, a, s), \mu', s') \} \\ & \leq \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta E \widetilde{W}(\ell(x, a, s), \mu', s') \}\end{aligned}$$

It follows that

$$\begin{aligned}& \min_{\gamma \geq 0} \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta E \widehat{W}(\ell(x, a, s), \varphi(\mu, \gamma), s') \} \\ & \leq \min_{\gamma \geq 0} \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta E \widetilde{W}(\ell(x, a, s), \varphi(\mu, \gamma), s') \}\end{aligned}$$

■

Notice that if  $W \in \mathcal{M}_{bc}$  and  $r \in \mathcal{R}$ ,  $(W+r)(x, \mu, s) = \mu(\omega+r)(x, \mu, s) = \mu\omega(x, \mu, s) + r\|\mu\|$ .

**Lemma 8A (discounting)** Assume **A4** and **A7b**. For any  $W \in \mathcal{M}_b$ , and  $r \in \mathcal{R}_+$ ,  $T^*(W+r) \leq T^*W + \beta r$ .

**Proof** First notice that, for any  $(x, \mu, s)$  and  $\gamma \geq 0$ ,

$$\begin{aligned}& \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta E(W+r)(\ell(x, a, s), \varphi(\mu, \gamma), s') \} \\ & = \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta EW(\ell(x, a, s), \varphi(\mu, \gamma), s') + \beta r \|\varphi(\mu, \gamma)\| \} \\ & = \max_{a \in A(x,s)} \{ \mu h_0(x, a, s) + \gamma h_1(x, a, s) + \beta EW(\ell(x, a, s), \varphi(\mu, \gamma), s') \} + \beta r \|\varphi(\mu, \gamma)\|,\end{aligned}$$

Now, given  $(x, \mu, s)$  and  $a \in A(x, s)$ , denote by  $\gamma^+(a)$  the solution to the following problem

$$\begin{aligned}& \min_{\{\gamma \geq 0: \|\gamma\| \leq B\|\mu\|\}} \left\{ \begin{array}{l} \mu h_0(x, a, s) + \beta E \sum_{j=0}^k \mu^j (\omega_j + r)(\ell(x, a, s), \varphi(\mu, \gamma), s') \\ + \sum_{j=0}^l \gamma^j [h_1^j(x, a, s) + \beta E (\omega_j + r)(\ell(x, a, s), \varphi(\mu, \gamma), s')] \end{array} \right\} \\ & = \mu h_0(x, a, s) + \beta E \sum_{j=0}^k \mu^j \omega_j(\ell(x, a, s), \varphi(\mu, \gamma^+(a)), s') \\ & + \gamma^+(a) [h_1(x, a, s) + \beta E \omega(\ell(x, a, s), \varphi(\mu, \gamma^+(a)), s')] + \beta r \|\varphi(\mu, \gamma^+(a))\|\end{aligned}$$

and let  $\gamma^*(a)$  be the solution to

$$\begin{aligned} & \min_{\{\gamma \geq 0: \|\gamma\| \leq B\|\mu\|\}} \left\{ \begin{array}{l} \mu h_0(x, a, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a, s), \varphi(\mu, \gamma), s') \\ \sum_{j=0}^l \gamma^j \left[ h_1^j(x, a, s) + \beta \mathbf{E} \omega_j(\ell(x, a, s), \varphi(\mu, \gamma), s') \right] \end{array} \right\} \\ &= \mu h_0(x, a, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a, s), \varphi(\mu, \gamma^*(a)), s') \\ &+ \gamma^*(a) [h_1(x, a, s) + \beta \mathbf{E} \omega(\ell(x, a, s), \varphi(\mu, \gamma^*(a)), s')]; \end{aligned}$$

therefore,

$$\begin{aligned} & \mu h_0(x, a, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a, s), \varphi(\mu, \gamma^+(a)), s') \\ &+ \gamma^+(a) [h_1(x, a, s) + \beta \mathbf{E} \omega(\ell(x, a, s), \varphi(\mu, \gamma^+(a)), s')] + \beta r \|\varphi(\mu, \gamma^+(a))\| \\ &\leq \mu h_0(x, a, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a, s), \varphi(\mu, \gamma^*(a)), s') \\ &+ \gamma^*(a) [h_1(x, a, s) + \beta \mathbf{E} \omega(\ell(x, a, s), \varphi(\mu, \gamma^*(a)), s')] + \beta r \|\varphi(\mu, \gamma^*(a))\|. \end{aligned}$$

Piecing things together (denoting, as usual, by  $a^* \equiv a^*(x, \mu, s)$  and  $\gamma^* \equiv \gamma^*(x, \mu, s)$ ); that is,  $\gamma^* \equiv \gamma^*(a^*)$ ), we have:

$$\begin{aligned} & T^*(W + r)(x, \mu, s) \\ &= \min_{\gamma \geq 0} \max_{a \in A(x, s)} \left\{ \begin{array}{l} \mu h_0(x, a, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a, s), \varphi(\mu, \gamma), s') \\ + \sum_{j=0}^l \gamma^j \left[ h_1^j(x, a, s) + \beta \mathbf{E} \omega_j(\ell(x, a, s), \varphi(\mu, \gamma), s') \right] \end{array} \right\} \\ &\leq \mu h_0(x, a^*, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a^*, s), \varphi(\mu, \gamma^*, s')) \\ &+ \gamma^* [h_1(x, a^*, s) + \beta \mathbf{E} \omega(\ell(x, a^*, s), \varphi(\mu, \gamma^*), s')] + \beta r \|\varphi(\mu, \gamma^*)\| \\ &= \mu h_0(x, a^*, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a^*, s), \varphi(\mu, \gamma^*, s')) + \beta r \|\varphi(\mu, \gamma^*)\| \\ &\leq \mu h_0(x, a^*, s) + \beta \mathbf{E} \sum_{j=0}^k \mu^j \omega_j(\ell(x, a^*, s), \varphi(\mu, \gamma^*, s')) + \beta r(1 + B) \|\mu\| \end{aligned}$$

By homogeneity, without loss of generality we can choose an arbitrary  $\mu \neq 0$ , such that:  $\|\mu\| \leq (1 + B)^{-1}$ . The above inequalities show that  $T^*(W + r) \leq T^*(W) + \beta r \blacksquare$

**Lemma 9A (Contraction property):** The argument is the standard Blackwell's argument. We show that the contraction property is satisfied. Let  $W, \widehat{W} \in \mathcal{M}_{bc}$ .

Notice that  $W \leq \widehat{W} + \|W - \widehat{W}\|$ , then using the results of Lemmas 7A and 8A,

$$T^*W \leq T^*(\widehat{W} + \|W - \widehat{W}\|) \leq T^*(\widehat{W}) + \beta\|W - \widehat{W}\|,$$

reversing the roles of  $W$  and  $\widehat{W}$ , we obtain that

$$\|T^*W - T^*\widehat{W}\| \leq \beta\|W - \widehat{W}\|;$$

therefore,  $T^*$  is a contraction mapping ■



## REFERENCES

- ABRAHAM, A. AND N. PAVONI (2005): "The Efficient Allocation of Consumption under Moral Hazard and Hidden Access to the Credit Market," *Journal of the European Economic Association*, vol. 3(2-3), 370-381.
- ABREU, D., D. PEARCE AND E. STACCHETTI (1990): "Toward a Theory of Discounted Repeated Games with Imperfect Monitoring," *Econometrica* 58, 1041-1063.
- ACEMOGLU, D., M. GOLOSOV AND A. TSYVINSKII (2011): "Power Fluctuations and Political Economy", *Journal of Economic Theory* (forthcoming)
- AIYAGARI, R., A. MARCET, T. J. SARGENT AND J. SEPPALA (2002): "Optimal Taxation without State-Contingent Debt", *Journal of Political Economy* 110, 1220-54.
- ANAGNOSTOPOULOS, A., E. CARCELES-POVEDA AND A. MARCET (2011) "Equity Financing", working paper.
- ATTANASIO, O. AND J.-V. RIOS-RULL (2000): "Consumption Smoothing in Island Economies: Can Public Insurance Reduce Welfare?," *European Economic Review* 44, 1225-58.
- BEWLEY, T. F. (1972): "Existence of Equilibria in Economies with Infinitely Many Commodities," *Journal of Economic Theory* 4: 514-540.
- BEWLEY, T. F. (1991): "A Theorem on the Existence of Competitive Equilibria in a Market with a Finite Number of Agents and Whose Commodity Space is  $\mathbf{L}_\infty$ ", in *Equilibrium Theory in Infinite Dimensional Spaces* ed. by M.A. Khan and N.C. Yannelis. Springer-Verlag.
- BROER, T. (2009): "Stationary Equilibrium Distributions under Limited Commitment," EUI Wp ECO 2009/39, European University Institute.
- CHANG, R. (1998): "Credible Monetary Policy with Long-Lived Agents: Recursive Approaches," *Journal of Economic Theory* 81, 431-461.
- COLE, H. AND F. KUBLER (2010): "Recursive Contracts, Lotteries and Weakly Concave Pareto Sets," mimeo, University of Pennsylvania.
- COOLEY, T. F. (ed.) (1995): *Frontiers of Business Cycle Research*. Princeton, N.J.: Princeton University Press.
- COOLEY, T. F, R. MARIMON AND V. QUADRINI (2004): "Optimal Financial Contracts with Limited Enforceability and the Business Cycle," *Journal of Political Economy* 112, 817-47.
- CRONSHAW, M. AND D. LUENBERGER (1994): "Strongly Symmetric Subgame Perfect Equilibria in Infinitely Repeated Games with Perfect Monitoring and Discounting," *Games and Economic Behavior*, 6, 220-237.

- DEBORTOLI, D. AND R. NUNES, (2010). "Fiscal policy under loose commitment," *Journal of Economic Theory*, vol. 145(3), 1005-1032, May.
- DUNFORD, N. AND J. T. SCHWARTZ (1957): *Linear Operators, Part I*. New York: John Wiley & Sons.
- EPPLE, D., L. HANSEN AND W. ROBERDS (1985): "Quadratic Duopoly Models of Resource Depletion," in *Energy, Foresight, and Strategy*, edited by T. J. Sargent, Resources for the Future.
- FERRERO, G. AND A. MARCET (2005): "Limited Commitment and Temporary Punishment", Universitat Pompeu Fabra Ph. D. thesis (Chapter 2).
- GREEN, E. J. (1987): "Lending and the Smoothing of Uninsurable Income" in *Contractual Arrangements for Intertemporal Trade* ed. by E. C. Prescott and N. Wallace, University of Minnesota Press.
- KEHOE, P. AND F. PERRI (2002): "International Business Cycles with Endogenous Market Incompleteness," *Econometrica* 70(3), 907-928.
- KOCHERLAKOTA, N. R. (1996): "Implications of Efficient Risk Sharing without Commitment," *Review of Economic Studies* 63 (4), 595-609.
- KYDLAND, F. E. AND E. C. PRESCOTT (1977): "Rules Rather than Discretion: The Inconsistency of Optimal Plans," *Journal of Political Economy* 85, 473-92.
- KYDLAND, F. E. AND E. C. PRESCOTT (1980): "Dynamic Optimal Taxation, Rational Expectations and Optimal Control," *Journal of Dynamics and Control* 2, 79-91.
- LEVINE, P. AND D. CURRIE (1987): "The design of feedback rules in linear stochastic rational expectations models," *Journal of Economic Dynamics and Control*, 11(1), 1-28.
- LJUNGQVIST L. AND T. SARGENT (2004): *Recursive Macroeconomic Theory*. The MIT Press.
- LUCAS, R. E. AND N. STOKEY (1983): "Optimal Fiscal and Monetary Policy," *Journal of Monetary Economics*.
- LUENBERGER, D. G. (1969): *Optimization by Vector Space Methods*. New York: Wiley.
- MELE A. (2010): "Repeated Moral Hazard and Recursive Lagrangeans," , Economics Department wp., Oxford University.
- MARCET, A. (2008): "Recursive Contracts," in *The New Palgrave Dictionary* ed. by S.N. Durlauf and L.E. Blume.

- MARCET, A. AND R. MARIMON (1992): "Communication, Commitment and Growth," *Journal of Economic Theory* 58:2, 219-249.
- MARCET, A. AND R. MARIMON (1998, 1999): "Recursive Contracts " *European University Institute, ECO 1998 #37 WP , Universitat Pompeu Fabra WP # 337*.
- MARIMON, R., MESSNER M. AND N. PAVONI (2011): "Solving Recursive Contracts with Non-unique Solutions," mimeo, European University Institute.
- MARIMON, R. AND P. RENDAHL (2011): "Recursive Bargaining with Endogenous Participation ", mimeo, University of California at Davis.
- MARIMON, R. AND V. QUADRINI (2011): "Competition, Human Capital and Income Inequality with Limited Commitment", *Journal of Economic Theory* (forthcoming).
- MESSNER M. AND N. PAVONI (2004): "On the Recursive Saddle Point Method," IGER Working paper 255, Universita Bocconi.
- PHELAN, C. AND E. STACCHETTI (2001): "Sequential Equilibria in a Ramsey Tax Model," *Econometrica* 69, 1491-518.
- ROCKAFELLAR, R.T. (1970): *Convex Analysis*. Princeton, NJ: Princeton University Press..
- ROCKAFELLAR, R.T. (1981): *The Theory of Subgradients and its Applications to Problems of Optimization. Convex and Nonconvex Functions*. Berlin: Heldermann Verlag..
- SARGENT, T.J.(1987): *Macroeconomic Theory*, (Chapter XV), New York: Academic Press, Second edition.
- SLEET, C. AND S. YELTEKIN (2010): "The Recursive Lagrangian Method: Discrete Time," mimeo, Carengie-Mellon University.
- STOKEY, N. L., R. E. LUCAS AND E. C. PRESCOTT (1989): *Recursive Methods in Economic Dynamics*. Cambridge, Ma.: Harvard University Press.
- SVENSSON, L. AND N. WILLIAMS (2008): "Optimal Monetary Policy under Uncertainty: A Markov Jump-Linear-Quadratic-Approach," *Federal Reserve Bank of St. Louis Review*.
- THOMAS, J. AND T. WORRAL (1990): "Income Fluctuations and Asymmetric Information: An Example of Repeated Principal-Agent Problem," *Journal of Economic Theory* 51: 367-390.

**CENTRE FOR ECONOMIC PERFORMANCE**  
**Recent Discussion Papers**

1054	Olivier Cadot Leonardo Iacovone Denisse Pierola Ferdinand Rauch	Success and Failure of African Exporters
1053	Björn Eriksson Tobias Karlsson Tim Leunig Maria Stanfors	Gender, Productivity and the Nature of Work and Pay: Evidence from the Late Nineteenth-Century Tobacco Industry
1052	Hartmut Lehmann Jonathan Wadsworth	The Impact of Chernobyl on Health and Labour Market Performance
1051	Jörn-Steffen Pischke	Money and Happiness: Evidence from the Industry Wage Structure
1050	Tim Leunig Joachim Voth	Spinning Welfare: the Gains from Process Innovation in Cotton and Car Production
1049	Francesca Cornaglia Andrew Leigh	Crime and Mental Wellbeing
1048	Gianluca Benigno Hande Küçük-Tüger	Portfolio Allocation and International Risk Sharing
1047	Guy Mayraz	Priors and Desires: A Model of Payoff-Dependent Beliefs
1046	Petri Böckerman Alex Bryson Pekka Ilmakunnas	Does High Involvement Management Lead to Higher Pay?
1045	Christos Genakos Tommaso Valletti	Seesaw in the Air: Interconnection Regulation and the Structure of Mobile Tariffs
1044	Giordano Mion Luca David Oromolla	Managers' Mobility, Trade Status and Wages
1043	Javier Ortega Gregory Verdugo	Immigration and the Occupational Choice of Natives: A Factor Proportions Approach
1042	Nicholas Bloom Benn Eifert Aprajit Mahajan David McKenzie John Roberts	Does Management Matter? Evidence from India
1041	Joshua D. Angrist Stacey H. Chen Brigham R. Frandsen	Did Vietnam Veterans Get Sicker in the 1990s? The Complicated Effects of Military Service on Self-Reported Health

1040	Tanvi Desai Felix Ritchie	Effective Researcher Management
1039	Ralf Martin Mirabelle Muûls Laure B. de Preux Ulrich J. Wagner	Anatomy of a Paradox: Management Practices, Organisational Structure and Energy Efficiency
1038	Giordano Mion Linke Zhu	Import Competition from and Outsourcing to China: A Curse or Blessing for Firms?
1037	William Brown David Marsden	Individualisation and Growing Diversity of Employment Relationships
1036	John Van Reenen	Does Competition Raise Productivity through Improving Management Quality?
1035	Florence Kondylis Marco Manacorda	School Proximity and Child Labor Evidence from Rural Tanzania
1034	Lars Boerner Albrecht Ritschl	Communal Responsibility and the Coexistence of Money and Credit under Anonymous Matching
1033	Gianluca Benigno Pierpaolo Benigno Salvatore Nisticó	Second-Order Approximation of Dynamic Models with Time-Varying Risk
1032	Gianluca Benigno Chuigang Chen Christopher Otrok Alessandro Rebucci Eric R. Young	Financial Crises and Macro-Prudential Policies
1031	Dennis Novy	International Trade without CES: Estimating Translog Gravity
1030	Alex Bryson John Forth	The Evolution of the Modern Worker: Attitudes to Work
1029	Fabrice Defever	The Spatial Organization of Multinational Firms
1028	Elhanan Helpman Oleg Itskhoki Stephen Redding	Trade and Labor Market Outcomes
1027	Nicholas Oulton	Long Term Implications of the ICT Revolution: Applying the Lessons of Growth Theory and Growth Accounting

**The Centre for Economic Performance Publications Unit**

**Tel 020 7955 7284 Fax 020 7955 7595**

**Email [info@cep.lse.ac.uk](mailto:info@cep.lse.ac.uk) Web site <http://cep.lse.ac.uk>**